

AD-A067 752

MINNESOTA UNIV MINNEAPOLIS DEPT OF PSYCHOLOGY

F/G 9/2

COMPUTER PROGRAMS FOR SCORING TEST DATA WITH ITEM CHARACTERISTI--ETC(U)

FEB 79 I I BEJAR, D J WEISS

N00014-76-C-0627

UNCLASSIFIED


RR-79-1

NL

1 of 1  
ADA  
067752



END  
DATE  
FILMED  
6-79  
DDC



BDC FILE COPY,

ADA06752

CO  
SC  
IT  
Is  
an  
Da



**LEVEL II**




AD A067752

COMPUTER PROGRAMS FOR  
SCORING TEST DATA WITH  
ITEM CHARACTERISTIC CURVE MODEL

Isaac I. Bejar  
and  
David J. Weiss

DDC FILE COPY

RESEARCH REPORT 7  
FEBRUARY 1



PSYCHOMETRIC METHODS PROG  
DEPARTMENT OF PSYCHOL  
UNIVERSITY OF MINNES  
MINNEAPOLIS, MN 55

This research was supported by funds from the Air Force Human Resources Laboratory, Defense Advanced Research Projects Agency, Navy Personnel Research and Development Center, Army Research Institute, and Office of Naval Research, monitored by the Office of Naval Research.

Approved for public release; distribution unlimited.  
Reproduction in whole or in part is permitted for any purpose of the United States Government.

79 04 20 00

## **DISCLAIMER NOTICE**

**THIS DOCUMENT IS BEST QUALITY  
PRACTICABLE. THE COPY FURNISHED  
TO DDC CONTAINED A SIGNIFICANT  
NUMBER OF PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.**

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

9. REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Research Report, 79-1	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Computer Programs for Scoring Test Data with Item Characteristic Curve Models.		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Isaac I. Bejar and David J. Weiss	8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0627	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Psychology University of Minnesota Minneapolis, MN 55455	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS P.E.: 61153N PROJ. RR042.04 T.A.: RR042.04.01 W.U.: NRI50-389	
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research Arlington, VA 22217	12. REPORT DATE February 1979	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) RR-79-1 (12) 74p.	13. NUMBER OF PAGES 84	15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This research was supported by funds from the Air Force Human Resources Laboratory, Defense Advanced Research Projects Agency, Navy Personnel Research and Development Center, Army Research Institute, and the Office of Naval Research, and monitored by the Office of Naval Research.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) item characteristic curve theory    test theory    psychological testing ICC    test scoring    adaptive testing item response theory    ability testing    computerized testing latent trait theory    achievement testing    tailored testing		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Three computer programs are described for scoring test response data using item characteristic curve (ICC) or latent trait models. The rationale and mathematical basis of both maximum likelihood and Bayesian ICC scoring methods are presented, as well as some data comparing the two methods of scoring. The three computer programs are designed for scoring conventional (linear) test data (LINDSCO) in dichotomous response format, adaptive test dichotomous data (ADADSCO), and conventional (linear) test data scored by polychotomous ICC models (LINPSCO). Options available in these three general		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

Unclassified


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

406 024 79 04 20 002



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

purpose programs are described, and examples of the input and output are given for each program. Complete FORTRAN listings of the three programs are included. 

**LEVEL II**

ACCESSION NO.	
DTIC	White Section <input checked="" type="checkbox"/>
DDC	Dist Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	B

DDC  
RECEIVED  
APR 23 1979  
D

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## CONTENTS

Introduction .....	1
An Introduction to Test Scoring .....	1
Inadequacies of the Number-Correct Score .....	1
ICC-Based Scoring .....	2
Maximum Likelihood Scoring .....	2
Bayesian Scoring .....	5
Differences Among Scoring Methods .....	7
General Description of the Programs .....	9
Numerical Procedures .....	10
Dichotomous Data .....	10
Maximum Likelihood .....	10
Formulas for Derivatives .....	11
Computation of Information .....	13
Standard Error .....	14
Bayesian Scoring .....	14
Computation of Expected Proportion of Correct Answers .....	15
Polychotomous Data .....	16
Graded Models .....	16
Nominal Logistic Model .....	20
Computation of Information .....	20
Use of the Programs .....	21
Input .....	21
Program Parameters .....	24
LINDSCO (Card Set 2-10) .....	24
ADADSCO (Card Set 2-5) .....	25
LINPSCO (Card Set 2-7) .....	25
Item Pool .....	26
LINDSCO and ADADSCO .....	26
Editing of Item Parameter Estimates .....	26
LINPSCO .....	27
Test Response Data .....	28
LINDSCO .....	28
ADADSCO .....	28
LINPSCO .....	29
Output .....	29
Program Parameters .....	29
Item Parameters .....	29
LINDSCO and LINPSCO .....	29
ADADSCO .....	30
Computational Messages .....	30
Testee Data .....	30
LINDSCO .....	30
ADADSCO .....	31
LINPSCO .....	31
Availability .....	31
References .....	32

## Appendices

Appendix A:	Item Parameter Estimation Programs .....	34
Appendix B:	Example of Program Use .....	35
Appendix C:	LINDSCO Fortran Program Listing .....	54
Appendix D:	ADADSCO Fortran Program Listing .....	64
Appendix E:	LINPSCO Fortran Program Listing .....	73

### Acknowledgments

The computer programs described in this report were written by Sinan N. Neftci, Meera Mahesh Pondicherry, and Ann Marie Kohler. Their contributions are gratefully acknowledged. The assistance of James B. Sympson in preparing the section entitled ICC-Based Scoring is also appreciated.

### Disclaimer

While every attempt has been made to insure the accuracy of the computer programs described in this report, the authors assume no responsibility for the accuracy or performance of the programs.

Technical Editor: Barbara Leslie Camm



## COMPUTER PROGRAMS FOR SCORING TEST DATA WITH ITEM CHARACTERISTIC CURVE MODELS

Although latent trait test theory, or item characteristic curve (ICC) theory, has been developing since Lawley's (1943) paper more than 30 years ago, applications of the theory have appeared only recently. However, there are indications that latent trait test theory is beginning to reach the practitioner who is concerned with test development and usage in applied settings. This is evidenced, not only by the increasing number of journal articles concerned with latent trait test theory (e.g., the summer 1977 special issue of the Journal of Educational Measurement on applications of latent trait models) and in presentations and training sessions at professional meetings, but also by its application in adaptive (Weiss, 1976) or tailored (Lord, 1970) testing.

A potential disadvantage of latent trait test theory is that its use often involves complex computational procedures. To apply ICC models to the development of tests and their scoring, the psychometrician must be able to estimate the ICC parameters of the items in the test, and then use them in conjunction with the response data of a new group of testees in order to estimate their trait scores (e.g., ability or achievement levels). A number of computer programs are available for estimating ICC item parameters (these are summarized in Appendix Table A). However, there appeared to be no general programs available for scoring test data with ICC models when item parameter estimates were available from previous data sets. This report describes several programs designed to meet this need.

### An Introduction to Test Scoring

The problem of test scoring can be conceptualized as the process of summarizing a testee's answers to a set of test questions into a single number in such a way that the score will be indicative of the testee's position on the trait being measured by the test. The most common test scoring strategy is to add the number of correct answers and to transform the score into some type of standard score or percentile to add interpretability. Historically, the number-correct score has been used because it is easy to calculate, and in pre-computer days this was an essential requirement of a test scoring procedure. As a general procedure for scoring tests of ability and achievement, however, the number-correct score has several deficiencies.

### Inadequacies of the Number-Correct Score

One major problem with the number-correct score is that it is possible for the same number-correct score to be obtained in several different ways; that is, several response patterns can result in the same number-correct score. If the items in a test are all of equal difficulty and discrimination, and therefore are essentially replicates of each other, this will have little effect on the number-correct score, since different response patterns among

replicate items are of little consequence. But it is a very rare test--and one which would have little general measurement utility--which would have items that are all replicates of each other with regard to difficulty and discrimination.

When test items differ with respect to difficulty or discrimination, they are no longer replicates. Under these circumstances, different patterns of response to the same set of items convey different information with regard to a testee's trait level. The testee who correctly answers only five very difficult items in a test is likely of higher ability than the testee who correctly answers only five very easy items in the same test. Although the total number-correct score is the same for these two testees, their trait level estimates derived from latent trait or ICC theory will differ. An additional unattractive feature of the number-correct score is the fact that the number of possible scores is determined by the number of items in the test. Thus, if a test consists of only 10 items, only 10 unique scores are possible. Although this may be sufficient in some applications, in others it might be desirable to obtain a finer gradation of scores.

The inadequacy of the number-correct score as a general test-scoring procedure is most obvious when considering how to score responses of testees who have been administered different sets of items, as in adaptive or tailored testing. In these kinds of tests, number-correct scores are completely inappropriate, since different testees will receive items of different difficulties and discriminations as well as different numbers of items in an adaptive test. In addition, the proportion of correct responses obtained by all testees will be approximately the same in a well-designed adaptive test (e.g., Weiss, 1975).

#### ICC-Based Scoring

The scoring programs described in this report use considerably more refined approaches than a mere adding of correct answers and are usable for scoring both conventional and adaptive test data. This refinement is possible because ICC theory makes very explicit specifications about the relationship between performance on a test item and the testee's position on the trait,  $\theta$ . This relationship is referred to as the *item characteristic curve* (ICC; Lord & Novick, 1968) when the items are scored into two categories (correct or incorrect) or, when there are more than two score categories, as the *operating characteristic function* (Samejima, 1969).

In the context of latent trait test theory, scoring may be conceptualized as finding the value of  $\theta$  (i.e., the trait being measured) most "compatible," in some sense, with a given pattern of responses to the test items, given the ICC item parameters for each item answered. For maximum likelihood scoring, the score associated with a given response vector is that value of  $\theta$  for which the likelihood of the response vector is maximum. For Bayesian scoring, the score is usually either the value of  $\theta$  that minimizes the mean squared difference between the estimated  $\theta$  and the "true"  $\theta$ , or the value of  $\theta$  that is most probable given the observed responses.


Maximum likelihood scoring. The details of the maximum likelihood scoring procedure are presented below. However, a conceptual explanation based on two



dichotomously scored test items will serve to explicate its rationale.

Figure 1 shows response probability curves for two test items--Item 1, which was answered correctly (resulting in an ICC plot of the probability of a correct response), and Item 2, which was answered incorrectly (resulting in a descending plot of the probability of an incorrect response, or 1 minus the ICC). The ICC curves for the two items are described by three parameters: (1) difficulty,  $b$ , which is the location of the ICC on the trait ( $\theta$ ) continuum at the point of maximum slope of the ICC ( $b=-.5$  for Item 1 and  $.75$  for Item 2); (2) their discrimination,  $a$ , which is proportional to the slope of the ICC at  $b$  ( $a=.8$  for Item 1 and  $1.4$  for Item 2); and (3) "guessing,"  $c$ , the lower asymptote of the probability of a correct response at  $\theta=-\infty$  ( $c=.16$  for Item 1 and  $1-.78=.22$  for Item 2).

Figure 1  
Response Probability Plots for a Correctly Answered Item (Item 1)  
and an Incorrectly Answered Item (Item 2)



The first step in maximum likelihood scoring consists of determining the likelihood of the response pattern (correct response to Item 1 and incorrect response to Item 2). Assuming local independence, which means that responses to the test items have nothing in common except their relationship to the underlying trait,  $\theta$ , the likelihood of a response pattern at any value of  $\theta$  can be determined by multiplying the separate probabilities of the responses in the response pattern for that value of  $\theta$ . The value of  $\theta$  for which the likelihood is maximum is the maximum likelihood estimate of  $\theta$ .

Conceptually, this can be illustrated with the ICCs in Figure 1 by using discrete values of  $\theta$ , such as those shown in Table 1. For example, at  $\theta=-1.0$ , the probability of a correct response to Item 1 (scored as 1) is .442 and the

probability of an incorrect response to Item 2 (scored as 0) is .768; multiplying these values gives the likelihood of the [1,0] response pattern as .340. At  $\theta=+1.0$ , the probability of a correct response [1] to Item 1 is .903 and the probability of an incorrect response to Item 2 is .277; the likelihood of the [1,0] response pattern is therefore .250. Similarly, at  $\theta=0.0$ , the probability of a correct response to Item 1 is .718 and the probability of an incorrect response to Item 2 is .668, resulting in a likelihood for the [1,0] response pattern of .479. This process of computing likelihoods for the [1,0] response pattern can be repeated for a large number of values along the  $\theta$  continuum.

Table 1  
Probability of a Correct Response to  
Item 1 [ $P_1(\theta)$ ] and Probability of an  
Incorrect Response to Item 2 [ $Q_2(\theta)$ ] for  
Selected Values of  $\theta$  (Item 1:  $a=.8$ ,  
 $b=-.5$ ,  $c=.16$ ; Item 2:  $a=1.4$ ,  $b=.75$ ,  
 $c=.22$ ), and Values of the Likelihood  
Function [ $L(\theta)$ ]


$\theta$	$P_1(\theta)$	$Q_2(\theta)$	$L(\theta)$
-3.0	.187	.780	.146
-2.5	.212	.780	.165
-2.0	.257	.779	.200
-1.5	.332	.776	.257
-1.0	.442	.768	.340
.5	.580	.742	.430
0.0	.718	.668	.479
.5	.828	.503	.416
1.0	.903	.277	.250
1.5	.948	.112	.106
2.0	.973	.038	.037
2.5	.986	.012	.012
3.0	.993	.004	.004

The result of computing likelihoods for all possible values of  $\theta$  based on the response pattern and the relevant ICCs can be a plot of the likelihood values as a function of  $\theta$ . This plot, shown as a solid curve in Figure 2, is called a likelihood function. As can be seen, the maximum of the likelihood function in Figure 2 occurs at about  $\theta=0.0$  (actually .01). Thus,  $\theta=.01$  can be considered the maximum likelihood estimate of  $\theta$  associated with the [1,0] response pattern, given the parameters of the ICCs for the items generating that response pattern. The maximum likelihood  $\theta$  estimate is thus the value of  $\theta$  which maximizes likelihood of the given response pattern for items with the specified ICCs.

The generalization of the scoring method for more than two items is straightforward. For each value of  $\theta$ , the likelihood would be determined by multiplying the response probabilities for the appropriate ICCs (based on the specified response pattern) across all items that have been answered. Thus, for  $n$  items,  $n$  probabilities would be multiplied at each value of  $\theta$  to obtain the likelihoods. The resultant likelihood values for all values of  $\theta$

could be plotted; and the maximum of the likelihood function would be used to identify the value of  $\theta$  that gives the observed response pattern the greatest probability of occurrence.

Figure 2  
Likelihood Function and Bayesian Posterior  
Function for the [1,0] Response Pattern



Maximum likelihood scores are intuitively appealing; at the same time, they have a number of optimal statistical characteristics, at least asymptotically (i.e., when large numbers of items are administered). Of special relevance is the fact that as the number of items in the response pattern increases, it can be shown (Kendall & Stuart, 1961) that maximum likelihood estimates have minimum variance; and the reciprocal of that variance is known as the information function of  $\theta$ . As a consequence, different scores (i.e.,  $\theta$  estimates) can have different degrees of accuracy as estimators of  $\theta$  (Birnbaum, 1968; Samejima, 1969).

Bayesian scoring. Although the numerical details of maximum likelihood and Bayesian scoring are substantially different, the two methods are conceptually very similar. Bayesian scores are based on the likelihood function modified by the prior probability density function of  $\theta$ . The prior probability density function describes the assumed distribution of  $\theta$  in the population of individuals to be tested.

To illustrate, call  $L(\theta)$  the likelihood of the response pattern for a given  $\theta$  value. Now call  $f(\theta)$  the prior probability density associated with that value of  $\theta$ . The modified likelihood, which may be called  $p(\theta)$  is then

$$p(\theta) = f(\theta)L(\theta) / \int [f(\theta)L(\theta)]d\theta. \quad [1]$$



Equation 1 is called the posterior probability density function. Just as the maximum likelihood score is the value of  $\theta$  for which  $L(\theta)$  is maximum, one kind of Bayesian score is the value of  $\theta$  for which  $p(\theta)$  is maximum. Such scores are called Bayes modal estimates by Samejima (1969) because they are based on the mode of the posterior density function. A different type of Bayesian estimate is based on the mean of the posterior density function. Owen's (1975) Bayesian scoring procedure, which will be described in detail below, is an example of this approach. In his procedure, the prior probability densities are provided by a normal density function. Other Bayesian scoring procedures are also available (Sympson, 1977).

These concepts can be illustrated using the likelihoods associated with the [1,0] response pattern discussed earlier. Table 2 shows for several values of  $\theta$  the probability of a correct response to the first item [ $P_1(\theta)$ ]; the probability of an incorrect response to the second item [ $Q_2(\theta)$ ]; the likelihood of the response pattern [ $L(\theta)$ ] (these first three columns correspond to the data in Table 1); the prior probability densities [ $f(\theta)$ ], which in this case are ordinates of a normal distribution with mean of zero and standard deviation of 1; and the posterior density function [ $p(\theta)$ ], computed using Equation 1. For these data

$$\int [f(\theta)L(\theta)]d\theta \approx .348. \quad [2]$$

The resulting posterior density function, [ $p(\theta)$ ], is shown as the dashed curve in Figure 2.

Table 2  
Response Probabilities [ $P_1(\theta)$ ,  $Q_2(\theta)$ ], Likelihoods [ $L(\theta)$ ],  
Weights [ $w(\theta)$ ], and Posterior Density Function [ $p(\theta)$ ] for a  
Two-Item Response Pattern

$\theta$	$P_1(\theta)$	$Q_2(\theta)$	$L(\theta)$	$f(\theta)$	$p(\theta)$
-3.0	.187	.780	.146	.004	.002
-2.5	.212	.780	.165	.018	.009
-2.0	.257	.779	.200	.054	.031
-1.5	.332	.776	.257	.130	.096
-1.0	.442	.768	.340	.242	.236
-0.5	.580	.742	.430	.352	.435
0.0	.718	.668	.479	.399	.549
0.5	.828	.503	.416	.352	.421
1.0	.903	.277	.250	.242	.174
1.5	.948	.112	.106	.130	.040
2.0	.973	.038	.037	.054	.006
2.5	.986	.012	.012	.018	.001
3.0	.993	.004	.004	.004	.000

The mode of the posterior density function in Figure 2 is located near  $\theta=0$ , so the Bayesian modal estimate and the maximum likelihood estimate are about the same for this data. The Bayesian  $\theta$  estimate based on the mean of the  $p(\theta)$  distribution is  $-.12$ . This  $\theta$  estimate does not coincide with the maximum likelihood estimate ( $\hat{\theta}=.01$ ); as will be further shown below, estimates of  $\theta$  obtained from different ICC scoring methods do not generally agree.

#### Differences Among Scoring Methods

The programs described in this report are capable of scoring test data using most of the ICC response models available. The selection among models should not be arbitrary, especially when individual decisions are to be made on the basis of test scores. Dichotomous data can be scored by means of the one-, two-, and three-parameter ICC models, using either a normal or logistic ogive ICC. Thus, given the decision with regard to the number of parameters that describe the ICC, there still remains the problem of choosing between the normal or logistic ogive response models for scoring purposes. Unfortunately, there are as yet no firm guidelines for choosing between these two response models. Samejima (1969) has shown that the normal and logistic ogive models differ with respect to their scoring "philosophies," but the practical implications of these differences remain to be investigated.

To illustrate the differences among the models and different ICC scoring procedures, all response patterns for a five-item test were scored by maximum likelihood, assuming both normal and logistic ogive ICCs, and by Owen's (1975) Bayesian scoring method. Table 3 gives the item parameters assumed for the hypothetical five-item test. For all items, the  $c$  (guessing) parameter was set at 0.0, indicating that a two-parameter ICC model was used. Items varied in difficulty ( $b$ ) from  $-2$  to  $+2$  and had discriminations of 1.00 or 1.50.

Table 3  
Item Parameters for Five-Item Test

Item	$a$	$b$	$c$
1	1.00	-2.00	.00
2	1.50	-1.00	.00
3	1.00	0.00	.00
4	1.50	1.00	.00
5	1.00	2.00	.00

In a five-item test in which each item is scored dichotomously, there are  $2^5=32$  different response patterns. These response patterns are shown in Table 4 along with the scores associated with them. It is obvious from the data in Table 4 that for a given response pattern, the scores (all of which are on the same metric) differed somewhat. This indicates that the scoring procedures are not interchangeable.

For example, consider the five response patterns which have 20% correct, namely Patterns 2, 3, 5, 9, and 17. Not only do the  $\theta$  estimates (scores) for a given response pattern differ among the three scoring procedures, but there are some differences in the ordering of the  $\theta$  estimates derived from these response patterns within each procedure. For maximum likelihood scoring using

a normal ogive ICC, the ordering of the  $\theta$  estimates derived from the five response patterns was exactly the same as that obtained from the Bayesian scoring procedure, although the numerical values of the  $\theta$  estimates were uniformly higher for the Bayesian procedure. For both these scoring methods, there was a tendency for higher ability estimates to be obtained when a more difficult item was answered correctly. For example, the lowest  $\theta$  estimate was obtained by both scoring methods when the easiest item (Item 1) was answered correctly (Response Pattern 17); when only Item 2 was answered correctly (Response Pattern 9), the  $\theta$  estimates from both the Bayesian and maximum likelihood normal procedures increased. In addition, both scoring methods took into account the discriminations of the items involved. For example, Response

Table 4  
Scores Given to Each Response Pattern by Three Scoring Methods

Response Pattern	Maximum Likelihood		Bayesian
	Normal	Logistic	
1. 00000	$\infty^*$	$\infty^*$	-1.72
2. 00001	-.93	-1.60	-.64
3. 00010	-.61	-1.19	-.38
4. 00011	-.13	-.46	.11
5. 00100	-1.42	-1.60	-1.06
6. 00101	-.50	-.84	-.28
7. 00110	-.30	-.46	-.11
8. 00111	.13	.46	.30
9. 01000	-1.24	-1.19	-.89
10. 01001	-.23	-.46	-.15
11. 01010	.03	.00	.00
12. 01011	.50	.84	.41
13. 01100	-.60	-.46	-.42
14. 01101	.23	.46	.17
15. 01110	.39	.84	.28
16. 01111	.93	1.60	.64
17. 10000	-1.63	-1.60	-1.16
18. 10001	-.39	-.84	-.24
19. 10010	-.17	-.46	-.06
20. 10011	.30	.46	.39
21. 10100	-.78	-.84	-.58
22. 10101	.03	.00	.11
23. 10110	.17	.46	.23
24. 10111	.61	1.19	.62
25. 11000	-.42	-.46	-.29
26. 11001	.60	.46	.51
27. 11010	.78	.84	.63
28. 11011	1.42	1.60	1.09
29. 11100	.42	.46	.31
30. 11101	1.24	1.19	.93
31. 11110	1.63	1.60	1.08
32. 11111	$\infty^*$	$\infty^*$	1.55

\* For maximum likelihood scoring, it is not possible to score response patterns with all correct or incorrect answers.



Pattern 2 (with a correct response to Item 5, the most difficult item) was assigned higher scores than Pattern 5; but Pattern 2 was assigned lower scores than Pattern 3 (which had a correct response to Item 4, the second most difficult item), since in Response Pattern 3 a correct answer was given to an item (Item 4) with a higher discrimination than that of Pattern 2 (Item 5).

On the other hand, assuming a logistic ogive ICC for the maximum likelihood scoring procedure, estimated values of  $\theta$  were related to the discriminations of the items answered correctly. Those response patterns for which the discriminations of the items answered correctly were the same were assigned the same score, namely -1.60 for Patterns 2, 5, and 17 and -1.19 for Patterns 3 and 9. For the latter two response patterns, the discriminations of the items answered correctly were 1.50; for the former three response patterns, they were 1.00. Thus, the magnitude of the scores was a function of the item discriminations, and the item difficulties did not affect the  $\theta$  estimates.

These data indicate that the assumption of different forms of the ICC within the maximum likelihood scoring procedure will, in general, result in different  $\theta$  estimates. Since the Bayesian  $\theta$  estimates were different from both the maximum likelihood estimates, these three ICC-based scoring procedures are not interchangeable. However, additional research is required to further delineate the similarities and differences among the  $\theta$  estimates derived by different ICC-based scoring procedures and, more importantly, to assess the implications of these differences in practical applications.

#### General Description of the Programs

This report describes three computer programs for scoring test data with ICC models--LINDSCO, ADADSCO, and LINPSCO. Table 5 summarizes the major features of these programs. LINDSCO (LInear Dichotomous SCOring) is designed

Table 5  
Summary of Program Capabilities

Model and Scoring Procedure	Dichotomous		Polychotomous (LINPSCO)	
	Linear (LINDSCO)	Adaptive (ADADSCO)	Graded	Nominal
Logistic Ogive				
Bayesian <sup>a</sup>	NO	NO	NO	NO
Maximum Likelihood	YES	YES	YES	YES
Normal Ogive				
Bayesian <sup>a</sup>	YES	YES	NO	NO
Maximum Likelihood	YES	YES	YES	NO

<sup>a</sup>The Bayesian scoring procedure is based on Owen (1975).

to be used for scoring test data for conventional (linear) tests in which all items are administered to each testee. It requires responses to be dichotomous; that is, responses are scored into one of two categories, such as "correct"

and "incorrect." Omissions are permitted, but they are ignored in the computations. The number of omitted items is tallied from the number of items administered and reported as part of the output for each testee. Either the normal or logistic ogive response model can be used with ICCs described by one, two, or three parameters for maximum likelihood scoring. Response patterns may also be scored by Owen's (1975) Bayesian method which assumes a normal ogive ICC. The user can also specify, in addition to a total test score, subscores on as many as 25 subscales.

ADADSCO (ADaptive Dichotomous SCOring) is similar to LINDSCO, but it is designed specifically for scoring item response data derived from adaptive testing. Since in adaptive testing each respondent answers a different set of test items, the program must locate for each testee the item parameter estimates of each attempted item; LINDSCO, in contrast, does the item search only once. ADADSCO also differs from LINDSCO in that it has no subscale scoring capabilities.

LINPSCO (Linear Polychotomous SCOring) is designed to score data from linear (conventional) tests in which each testee is administered all items, and items are scored into more than two categories. Three models are available: the graded normal and logistic ogive models (Samejima, 1969), and the nominal logistic model (Bock, 1972). In LINPSCO only maximum likelihood scoring is available, and subscale scoring is not possible.

All three programs compute both test information and response pattern information values when maximum likelihood scoring is used. Response pattern information (Samejima, 1973) provides an estimate of the precision of measurement for a specified response pattern and can be used to compare the quality of trait estimates derived from specific test administration and/or scoring procedures (e.g., Bejar, Weiss, & Gialluca, 1977).

## NUMERICAL PROCEDURES

### Dichotomous Data

#### Maximum Likelihood

The numerical procedure for maximum likelihood scoring of dichotomous data consists of two stages. In the first stage an initial estimate is sought by the bisection method. Once this initial estimate is obtained, it is refined further by the Newton-Raphson method.

The bisection routine begins in the interval  $\pm 5.00$ . If the sign of the first derivative of the likelihood function during the first iteration is the same when evaluated at 5.00 and at -5.00, a value of 0.0 is returned as the initial estimate of  $\theta$ . Otherwise, five additional iterations are performed. After the sixth iteration, the width of the interval has been reduced to  $10/(2^6) = 10/64 = .15$ . The midpoint of that interval is the initial estimate which is then refined further by Newton-Raphson iterations of the form:

$$\hat{\theta}_{m+1} = \hat{\theta}_m - (f'/f'') \quad , \quad [3]$$



where

$\hat{\theta}_{m+1}$  is the new estimate,

$\hat{\theta}_m$  is the estimate from the last iteration,

$f'$  is the first derivative of the log-likelihood function evaluated at  $\hat{\theta}_m$ , and

$f''$  is the second derivative of the log-likelihood function evaluated at  $\hat{\theta}_m$ .

This iterative process is continued until  $|\hat{\theta}_{m+1} - \hat{\theta}_m| < .005$ . If that criterion has not been met at the end of 50 iterations, the case is said to be nonconvergent.

Formulas for derivatives. Let  $v = \{u_g, g=1, 2, \dots, n\}$  be a response vector such that

$$u_g = \begin{cases} 1 & \text{if the item is answered correctly} \\ 0 & \text{if the item is answered incorrectly.} \end{cases}$$

Note that for scoring purposes, the response vector does not include rejected or omitted items. The probability that  $u_g=1$  for a given value of  $\theta$  and item parameters  $a_g$ ,  $b_g$ , and  $c_g$  is given by

$$P_g(\theta) = c_g + (1-c_g)[1 + e^{-1.7a_g(\theta - b_g)}]^{-1} \quad [4]$$

for the logistic ogive model and by

$$\begin{aligned} P_g(\theta) &= c_g + (1-c_g) \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{a_g(\theta - b_g)} e^{-t^2/2} dt \\ &= c_g + (1-c_g) \Phi [a_g(\theta - b_g)] \end{aligned} \quad [5]$$

for the normal ogive model, where  $\Phi$  stands for the standard cumulative normal distribution.

The log-likelihood function for the response vector is

$$\begin{aligned} L_v(\theta) &= \sum_g \log P_g(\theta)^{u_g} Q_g(\theta)^{1-u_g} \\ &= \sum_g [u_g \log P_g(\theta) + (1-u_g) \log Q_g(\theta)] \end{aligned} \quad [6]$$

where  $Q_g(\theta) = 1 - P_g(\theta)$  and  $P_g(\theta)$  is given by either Equation 4 or 5.

In general, the first and second derivatives of the log-likelihood function of a response vector are given by

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \left\{ u_g \left[ \left( \frac{1}{P_g(\theta)} \right) \left( \frac{\partial P_g(\theta)}{\partial \theta} \right) \right] + (1-u_g) \left[ \left( \frac{1}{Q_g(\theta)} \right) \left( \frac{\partial Q_g(\theta)}{\partial \theta} \right) \right] \right\} \quad [7]$$

and

$$\begin{aligned} \frac{\partial^2 L_v(\theta)}{\partial \theta^2} = \sum_g \left\{ u_g \left[ \left( \frac{1}{P_g^2(\theta)} \right) \left( \frac{\partial P_g(\theta)}{\partial \theta} \right)^2 - \left( \frac{1}{P_g(\theta)} \right) \left( \frac{\partial^2 P_g(\theta)}{\partial \theta^2} \right) \right] \right. \\ \left. + (1-u_g) \left[ \left( \frac{1}{Q_g^2(\theta)} \right) \left( \frac{\partial Q_g(\theta)}{\partial \theta} \right)^2 + \left( \frac{1}{Q_g(\theta)} \right) \left( \frac{\partial^2 Q_g(\theta)}{\partial \theta^2} \right) \right] \right\} . \end{aligned} \quad [8]$$

For the logistic ogive model, after simplification and letting  $x = 1.7a_g(\theta - b_g)$ , these expressions are

$$\frac{\partial L_v(\theta)}{\partial \theta} = -1.7 \sum_g \left[ \frac{a_g e^x}{1+e^x} \right] + 1.7 \sum_g \left[ \frac{u_g a_g e^x}{c_g + e^x} \right] \quad [9]$$

and

$$\frac{\partial^2 L_v(\theta)}{\partial \theta^2} = -2.89 \sum_g \left[ \frac{a_g^2 e^x}{(1+e^x)^2} \right] + 2.89 \sum_g \left[ \frac{u_g a_g^2 c_g e^x}{(c_g + e^x)^2} \right] . \quad [10]$$

For the normal ogive model, letting  $x = -a_g^2(\theta - b_g)^2/2$ , the corresponding expressions are

$$\frac{L_v(\theta)}{\partial \theta} = \sum_g \left[ \frac{u_g (2\pi)^{-1/2} a_g (1-c_g) e^x}{c_g + (1-c_g) \Phi[a_g(\theta - b_g)]} - \frac{(1-u_g) (2\pi)^{-1/2} (1-c_g) a_g e^x}{1 - \{c_g + (1-c_g) \Phi[a_g(\theta - b_g)]\}} \right] \quad [11]$$

and

$$\begin{aligned} \frac{\partial^2 L_v(\theta)}{\partial \theta^2} = \sum_g \left\{ u_g \left[ - \frac{[(2\pi)^{-1/2}(1-c_g)a_g e^x]^2}{\{[c_g - (1-c_g)\Phi[a_g(\theta-b_g)]]\}^2} - \frac{(2\pi)^{-1/2}a_g^3(\theta-b_g)(1-c_g)e^x}{\{c_g - (1-c_g)\Phi[a_g(b_g-\theta)]\}} \right] + \right. \\ \left. \sum_g (1-u_g) \left[ - \frac{[(2\pi)^{-1/2}(1-c_g)a_g e^x]^2}{1-\{c_g + (1-c_g)\Phi[a_g(\theta-b_g)]\}^2} + \frac{(2\pi)^{-1/2}a_g^3(\theta-b_g)(1-c_g)e^x}{1-\{c_g + (1-c_g)\Phi[a_g(\theta-b_g)]\}} \right] \right\} \quad [12] \end{aligned}$$

*Computation of information.* With maximum likelihood scoring, two measures of information are computed for each response pattern. One is response pattern information (Samejima, 1973) denoted by  $\hat{I}(\hat{\theta})$ ; the other is test information (Birnbaum, 1968; Samejima, 1969) denoted by  $I(\hat{\theta})$ . Test information is defined as the expected value of the second derivative of the log-likelihood function, i.e.,

$$I(\hat{\theta}) = -\sum_g \left[ E \left\{ \frac{\partial^2 \log L_v(\theta)}{\partial \theta^2} \right\} \right] \quad [13]$$

Response pattern information, on the other hand, is defined by

$$\hat{I}(\hat{\theta}) = -\sum_g \left[ \frac{\partial^2 \log L_v(\theta)}{\partial \theta^2} \right] \quad [14]$$

that is, the "observed," as opposed to expected, value of the second derivative of the log-likelihood function evaluated at  $\hat{\theta}$ .

These two measures of information will be the same for models in which there is a sufficient statistic for the response vector. In particular, this is true in the one- and two-parameter logistic ogive models. It is also true for the "zero" parameter normal ogive model, i.e., when the items are parallel. The value of  $\hat{I}(\hat{\theta})$  for a given response pattern is simply the value of the second derivative of the log-likelihood function at the last iteration, i.e., evaluated at the estimated value of  $\theta$ .

$I(\hat{\theta})$  is computed by

$$I(\hat{\theta}) = \sum_g \frac{\{P'_g(\hat{\theta})\}^2}{P_g(\hat{\theta})\{1.0 - P_g(\hat{\theta})\}} \quad [15]$$



where  $P_g(\theta)$  is given, in general, by Equation 4 for the logistic ogive model and by Equation 5 for the normal ogive model.

For the normal ogive model,

$$P'_g(\hat{\theta}) = \frac{a_g(1-c_g)}{\sqrt{2\pi}} [e^{-a_g^2[\hat{\theta}-b_g]^2/2}] ; \quad [16]$$

and for the logistic ogive model,

$$P'_g(\hat{\theta}) = \frac{1.7a_g(1-c_g)e^{1.7a_g(\hat{\theta}-b_g)}}{[1+e^{1.7a_g(\hat{\theta}-b_g)}]^2} . \quad [17]$$

Standard error. The standard error of measurement associated with  $\hat{\theta}$  is computed as  $1/\sqrt{I(\hat{\theta})}$ , that is, the reciprocal square root of response pattern information evaluated at  $\hat{\theta}$ .

#### Bayesian Scoring

The Bayesian scoring procedure used by LINDSCO and ADADSCO is derived from Owen's (1975) sequential adaptive testing strategy. However, since the present application assumes that the test items have already been administered, only the scoring aspect is of interest.

The procedure makes the assumption that the prior distribution of  $\theta$  is normal, with mean  $\mu_0=0.0$  and variance  $\sigma_0^2=1.00$ , where subscript 0 denotes the fact that no items have yet been administered. After the  $m^{\text{th}}$  item is administered, the mean and variance of the posterior density function are computed according to the following equations. If the response to the  $m+1^{\text{th}}$  item is correct,

$$\mu_{m+1} = E(\theta|1) = \mu_m + (1-c_g) \left( \sqrt{\frac{\sigma_m^2}{\frac{1}{a_g^2} + \sigma_m^2}} \right) \left( \frac{\phi(D)}{c_g + (1-c_g)\phi(-D)} \right) \quad [18]$$

and

$$\sigma_{m+1}^2 = \text{var}(\theta|1) = \sigma_m^2 \left\{ 1 - \left( \frac{1-c_g}{1 + \frac{1}{a_g^2 \sigma_m^2}} \right) \left( \frac{\phi(D)}{A} \right) \left( \frac{(1-c_g)\phi(D)}{A} - D \right) \right\} . \quad [19]$$

Following an incorrect answer,

$$\mu_{m+1} = E(\theta|0) = \mu_m - \left( \frac{\sigma_m^2}{\sqrt{\frac{1}{\alpha_g^2} + \sigma_m^2}} \right) \left( \frac{\phi(D)}{\Phi(D)} \right) \quad [20]$$

and

$$\sigma_{m+1}^2 = \text{var}(\theta|0) = \sigma_m^2 \left\{ 1 - \left( \frac{\phi(D)}{1 + \frac{1}{\alpha_g^2 \sigma_m^2}} \right) \left( \frac{\frac{\phi(D)}{\Phi(D)} + D}{\Phi(D)} \right) \right\}. \quad [21]$$

In Equations 18 through 21 (from Owen, 1975),

$\phi(D)$  is the normal probability density function,  
 $\Phi(D)$  is the cumulative normal distribution function,

$$D = \frac{b_g - \mu_m}{\sqrt{\frac{1}{\alpha_g^2} + \sigma_m^2}}, \text{ and} \quad [22]$$

$$A = c_g + (1 - c_g) \Phi(-D). \quad [23]$$

After the last item has been administered, the posterior mean is the estimated  $\theta$  and the posterior variance is a measure of the error associated with that estimate. Because the posterior distribution after every item is administered is approximated by a normal distribution in this procedure, there is a certain amount of inaccuracy in the estimate. Moreover, the resulting scores are order dependent (Sympson, 1977), i.e., if a response vector were to be scored after rearranging the items, the resulting  $\theta$  estimate would be slightly different.

#### Computation of Expected Proportion of Correct Answers

The expected proportion of correct answers (EXPTOT) is defined as

$$\text{EXPTOT} = \sum_g P_g(\hat{\theta}) / NI, \quad [24]$$

where  $P_g(\hat{\theta})$  is computed from Equation 4 for the logistic ogive model and Equation 5 for normal ICCs.  $NI$  is the number of items on which the estimate of  $\theta$  is based. EXPTOT is simply an estimate of the true score associated with  $\hat{\theta}$  (Lord & Novick, 1968, p. 387).

### Polychotomous Data

LINPSO is capable of scoring polychotomous data when item parameters have been estimated according to a graded model of either normal or logistic ogive form (Samejima, 1969) or according to Bock's (1972) nominal logistic model. For the graded model, the numerical procedure consists of a bisection stage of six iterations followed by Newton-Raphson iterations. For the nominal logistic model, the initial estimate obtained from the bisection stage is refined further by the secant method rather than by Newton-Raphson iterations.

In each case, the bisection phase begins in the interval  $\pm 5.00$ . During the first iteration, if the sign of the first derivative of the log-likelihood function is the same when evaluated at 5.00 and at -5.00, a value of 0.0 is returned as the initial estimate. Otherwise, five additional iterations are performed. After six iterations, the width of the interval is reduced to  $10/(2^6) = 10/64 = .15$ . The midpoint of that interval is taken as the initial estimate.

The Newton-Raphson procedure used with the graded models refines the initial estimate with iterations of the form shown in Equation 3. This iterative procedure is continued until  $|\hat{\theta}_{m+1} - \hat{\theta}_m|$  is less than .005 or the number of iterations is greater than 50. The secant procedure is similar to Newton-Raphson iterations, except that  $f''$  in Equation 3 is an approximation to the second derivative of the log-likelihood function given by

$$f'' = \frac{f'(\hat{\theta}_m) - f'(\hat{\theta}_{m-1})}{(\hat{\theta}_m - \hat{\theta}_{m-1})} \quad [25]$$

### Graded Models

Let  $v = \{x_g, g=1, 2, \dots\}$  be a response vector exclusive of omitted and rejected items such that

$$x_g = \begin{cases} 1 & \text{if the "best" response was given} \\ 2 & \text{if the second "best" response was given} \\ \vdots & \\ m_g - 1 & \text{if the next to worst response was given} \\ m_g & \text{if the worst response was given.} \end{cases}$$

For the graded logistic ogive model, the probability that  $x_g$  takes one of the values between 1 and  $m_g$  is given in general by

$$P_{x_g}(\theta) = P_{x_g} = [1 + e^{y_{x_g}}]^{-1} - [1 + e^{y_{x_g-1}}]^{-1}, \quad [26]$$



where

$$y_{x_g} = -a_g D(\theta - b_{x_g}) \quad , \quad [27]$$

$$y_{x_g-1} = -a_g D(\theta - b_{x_g-1}) \quad , \quad \text{and} \quad [28]$$

$D = 1.7$  is a scaling factor.

When  $x_g = 1$ ,

$$P_{x_g} = [1 + e^{y_{x_g}}]^{-1} \quad . \quad [29]$$

When  $x_g = m_g$ ,

$$P_{x_g} = 1 - [1 + e^{y_{x_g-1}}]^{-1} \quad . \quad [30]$$

For the graded normal ogive model, the probability that  $x_g$  takes one of the values between 1 and  $m_g$  is given in general by

$$\begin{aligned} P_{x_g}(\theta) = P_{x_g} &= (2\pi)^{-1/2} \int_{y_{x_g-1}}^{y_{x_g}} e^{t^2/2} dt \\ &= \Phi[y_{x_g}] - \Phi[y_{x_g-1}] \quad , \end{aligned} \quad [31]$$

where

$$y_{x_g} = a_g(\theta - b_{x_g}) \quad [32]$$

and

$$y_{x_g-1} = a_g(\theta - b_{x_g-1}) \quad . \quad [33]$$

When  $x_g = 1$ ,

$$P_{x_g} = (2\pi)^{-1} \int_{-\infty}^{y_{x_g}} e^{t^2/2} dt \quad . \quad [34]$$

When  $x_g = m_g$ ,

$$P_{x_g} = 1 - (2\pi)^{-1} \int_{-\infty}^{y_{x_g}-1} e^{t^2/2} dt \quad [35]$$

The log-likelihood function for a given response vector is given by

$$\begin{aligned} L_v(\theta) &= \log \prod_g P_{x_g}^{r_{x_g}} \\ &= \sum_g r_{x_g} [\log P_{x_g}] , \end{aligned} \quad [36]$$

where

$$r_{x_g} = \begin{cases} 1 & \text{if the } x_g^{\text{th}} \text{ response category is chosen} \\ 0 & \text{otherwise} \end{cases} .$$

The general first derivative of the log-likelihood function is

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \sum_{x_g} r_{x_g} L_{x_g} \quad [37]$$

Samejima (1969) refers to  $L_{x_g}$  as the basic function. Since  $L_{x_g} = (\partial P_{x_g} / \partial \theta) (P_{x_g})^{-1}$

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \sum_{x_g} r_{x_g} \frac{\partial P_{x_g} / \partial \theta}{P_{x_g}} \quad [38]$$

The general second derivative of the log-likelihood function is given by

$$\frac{\partial^2 L_v(\theta)}{\partial \theta^2} = \sum_g \sum_{x_g} r_{x_g} \left[ -(L_{x_g})^2 + \frac{\partial^2 P_{x_g} / \partial \theta^2}{P_{x_g}} \right] \quad [39]$$

Specifically, for the graded logistic ogive model,

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \sum_{x_g} a_g 1.7 \{ 1 - P_{x_g}^* - P_{x_g}^{*-1} \} \quad [40]$$



and

$$\frac{\partial^2 L_v(\theta)}{\partial \theta^2} = \sum_g \sum_{x_g} 2.89 a_g^2 \frac{r_{x_g}}{P_{x_g}} \left[ - \left\{ (L_{x_g})^2 + (2Q_{x_g-1}^*) (P_{x_g}^* Q_{x_g}^*) - \right. \right. \\ \left. \left. (2Q_{x_g-1}^* - 1) (P_{x_g-1}^* Q_{x_g-1}^*) \right\} \right] , \quad [41]$$

where

$$P_{x_g}^* = [1 + e^{-1.7 a_g (\theta - b_{x_g})}]^{-1} , \quad [42]$$

$$P_{x_g-1}^* = [1 + e^{-1.7 a_g (\theta - b_{x_g-1})}]^{-1} , \quad [43]$$

$$P_o^* = 0 , \quad [44]$$

$$P_m^* = 1 , \quad [45]$$

$$Q_{x_g}^* = 1 - P_{x_g}^* , \text{ and } [46]$$

$$Q_{x_g-1}^* = 1 - P_{x_g-1}^* . \quad [47]$$

For the graded normal ogive model, letting  $z_{x_g} = -[a_g^2 (\theta - b_{x_g})^2] / 2$ , the corresponding expressions are

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \left[ \sum_{x_g} \left( \frac{r_{x_g} a_g}{\sqrt{2\pi}} [e^{z_{x_g}} - e^{z_{x_g-1}}] \right) / P_{x_g}(\theta) \right] \quad [48]$$

The second derivative is given by

$$\frac{\partial^2 L_v(\theta)}{\partial \theta^2} = \sum_g \left\{ \sum_{x_g} r_{x_g} \left[ - (L_{x_g})^2 \right] + \right. \\ \left. \left[ \frac{-a_g^3}{\sqrt{2\pi}} \{ (\theta - b_{x_g}) e^{z_{x_g}} - (\theta - b_{x_g-1}) e^{z_{x_g-1}} \} \right] / P_{x_g}(\theta) \right\} \quad [49]$$

When  $x_g = 1$ ,  $e^{z_{xg}-1} = 0$ , and  $P_{x_g}$  is given by Equation 34; when  $x_g = m_g$ ,  $e^{z_{xg}} = 0$ , and  $P_{x_g}$  is given by Equation 35.

### Nominal Logistic Model

For the nominal logistic model, the probability of  $x_g$ , given  $\theta$ , is given by

$$P_{x_g}(\theta) = P_{x_g} = e^{(\alpha_{x_g}\theta + \beta_{x_g})} / \sum_{s=1}^{m_g} e^{(\alpha_s\theta + \beta_s)}, \quad [50]$$

where  $\alpha_s$  and  $\beta_s$  are the slope and intercept parameter for the  $s^{\text{th}}$  response category.

The secant method requires only the first derivative of the log-likelihood function. That derivative is

$$\frac{\partial L_v(\theta)}{\partial \theta} = \sum_g \frac{\sum_{s=1}^{m_g} r_{x_g} (\alpha_{x_g} - \alpha_s) e^{\alpha_s\theta + \beta_s}}{\sum_{s=1}^{m_g} e^{\alpha_s\theta + \beta_s}}. \quad [51]$$

### Computation of Information

Response pattern information is computed as the value of the second derivative at the last iteration. For the nominal logistic model, that value is an approximation. Test information is computed from the general formula given by Samejima (1969),

$$I(\hat{\theta}) = \sum_g \sum_{x_g} (\partial P_{x_g} / \partial \theta)^2 P_{x_g}. \quad [52]$$

This expression involves only the first derivative of the response model. The appropriate expressions are listed below.

For the graded normal ogive model,

$$\frac{\partial P_{x_g}}{\partial \theta} = \frac{a_g}{\sqrt{2\pi}} \left[ e^{z_{xg}} - e^{z_{xg}-1} \right], \quad [53]$$

where  $z_{x_g} = -[\alpha_g^2(\theta - b_{x_g})^2] / 2$ .

For the graded logistic ogive model,

$$\frac{\partial P_{x_g}}{\partial \theta} = 1.7 \alpha_g [P_{x_g}^* (1 - P_{x_g}^*) - P_{x_g-1}^* (1 - P_{x_g-1}^*)], \quad [54]$$

where  $P_{x_g}^* = [1 + e^{-1.7\alpha (\theta - bx_g)}]^{-1}$ .

For the nominal logistic model,

$$\frac{\partial P_{x_g}}{\partial \theta} = \frac{[e^{(\alpha_{x_g}\theta + \beta_{x_g})} \sum_{s=1}^{m_g} e^{(\alpha_{s_g}\theta + \beta_{s_g})} (\alpha_{x_g} - \alpha_{s_g})]}{\sum_{s=1}^{m_g} e^{(\alpha_{s_g}\theta + \beta_{s_g})^2}} \quad [55]$$

#### USE OF THE PROGRAMS

##### Input

For each of the programs, three types of input are required:

1. The *Program Parameters*, which consist of specifications as to the number of items in the pool, the options chosen, the scoring key, and so forth.
2. The *Item Pool*, which contains the item parameter estimates on as many as 600 items for LINDSCO and ADADSCO, and 100 items for LINPSO.
3. The *Test Response Data* consists of testee name and identification number and each testee's item responses. For LINDSCO, item responses need not be dichotomized beforehand; for ADADSCO, they must be dichotomized unless a key is provided as part of the item pool. For ADADSCO, the number of items attempted and the identification number of each item attempted must also be supplied as part of the test response data. For LINPSO, the test response data must be supplied in such a way that the first category corresponds to the "best" response, while the last category corresponds to the "worst" response, based on previously obtained item parameterization data.

Testee response data containing all correct or incorrect answers cannot be scored by maximum likelihood. If such a response pattern is found, a message is printed, and the estimated  $\theta$  is set to 10.00 if all responses are correct and to -10.00 if all responses are incorrect. The information is set to 0.0 in both cases. Response patterns with all answers correct or incorrect present no problem for Bayesian scoring, and they are processed normally; however, a message is still printed. Appendix B gives examples of the use of each of these programs.



Table 6  
Input Program Parameters for LINDSCO, ADADSCO, and LINPSCO: Card Set 1

Columns	LINDSCO	ADADSCO	LINPSCO
1-4 (I4)	INUP, number of items in item pool. 600 is the maximum.	INUP, same as LINDSCO	Number of items in the pool. Maximum is 100.
5-8 (I4)	M, number of items in test. 300 is the maximum	MMAX, maximum number of items administered. 60 is the maximum.	M, number of items in the test. Maximum is 50.
9	blank	blank	blank
10 (II)	OPT1 1 = Punch the item parameter estimates corresponding to the items in the test.	OPT1 1 = Print the item parameter estimates corresponding to the items administered (this is done only for the first 10 testees).	OPT1 1 = Punch the item parameter estimates corresponding to the items in the test.
11 (II)	OPT2 1 = the item pool consists of M items, i.e., there will be no searching of items in the pool.	not used	OPT2 1 = the item pool consists of M items, i.e., there will be no searching of items in the pool.
12 (II)	OPT3 If 1, 2, or 3 item parameters will be edited; see "Editing of item parameters."	OPT3, same as LINDSCO	not used
13 (II)	OPT4, scoring algorithms and response model: 1 = maximum likelihood normal ogive 2 = maximum likelihood logistic 3 = Owen's Bayesian normal ogive	OPT4, same as LINDSCO	OPT4, response model: 1 = graded logistic ogive 2 = graded normal ogive 3 = nominal logistic

14-18 (F5.2)	TS, for Bayesian scoring. This is the prior mean of $\theta$ . Not used in maximum likelihood scoring.	TS, same as LINDSCO	D, scaling parameter for graded logistic model. If blank, will be set by default to 1.0; otherwise will usually be set by the user to 1.7.
19-23 (F5.2)	TSS, for Bayesian scoring. This is the prior standard deviation of $\theta$ . Not used in maximum likelihood scoring.	TSS, same as LINDSCO	not used
24-28 (F5.2)	AMAX, value of the $\alpha$ parameter. Used in editing. See "Editing of item parameters."	AMAX, same as LINDSCO	not used
29-33 (F5.2)	BMIN, lowest value of the $b$ parameter. Used in editing parameter estimates. See "Editing of item parameters."	BMIN, same as LINDSCO	not used
34-38 (F5.2)	BMAX, highest value of the $b$ parameter. Used in editing parameter estimates. See "Editing of item parameters."	BMAX, same as LINDSCO	not used
39-43 (F5.2)	CMAX, value of the $c$ parameter. Used in editing parameter estimates. See "Editing of item parameters."	CMAX, same as LINDSCO	not used
44-45 (I2)	blank	IFLAG, code for correct response	
46-47 (I2)	IOMIT, code for omitted response.	IOMIT, same as LINDSCO	IOMIT, same as LINDSCO
48-80	blank	blank	blank

### Program Parameters

Table 6 describes the input program parameters for all three programs, using Card Set 1 (all numeric information is right justified). After Card Set 1, the program parameter and input for each of the three programs differs, as indicated below.

#### LINDSCO (Card Set 2-10).

- |   |  |
|---|--|
| <u>Card Set 2 (8A10).</u>   | The variable format for the item pool is punched on this card, using I-fields (see Item Pool below).   |
| <u>Card Set 3 (16I5).</u>   | Punch in five-column fields the item identification number of the items in the test in the same order in which they appear in the test. Continue on as many cards as necessary.  |
| <u>Card Set 4 (80I1).</u>   | A "1" in a given column is punched to omit a specified item from all computations, e.g., if the 10th item is to be omitted, punch "1" in column 10; if the 100th item is to be omitted, punch "1" in column 20 of the second card. Continue on as many cards as necessary. |
| <u>Card Set 5 (80I1).</u>   | This card contains the scoring key for the test. In general, the $n^{\text{th}}$ column contains the key for the $n^{\text{th}}$ item, as in Card Set 4. Continue on as many cards as necessary.   |
| <u>Card Set 6 (8A10).</u>   | Variable format for reading the subject information and test response data (see Test Response Data below for field type specifications).   |
| <u>Card Set 7 (8A10).</u>   | The description of the run is written on three cards. The three cards must be included even if they are blank.   |
| <u>Card 8 (I5).</u>   | Punch the number of subscales to be scored in columns 1-5; maximum is 25. If no subscales are to be scored, punch "0" in column 5; in that case, this is the last card set.  |
| <u>Card Set 9 (2I5).</u><br>(Omit if the number of subscales is 0.)   | For each scale, punch the following information:<br>Columns 1-5: Number of items in subscale (maximum is 60).<br>Columns 6-10: Scale number. Repeat for each subscale beginning on a new card.   |
| <u>Card Set 10 (16I5).</u><br>(Omit if the number of subscales is 0.) | Punch in five-column fields the item identification number of the items in the subscales. Continue on as many cards as necessary. Repeat for each subscale, beginning on a new card for each subscale.   |



ADADSCO (Card Set 2-5).

Card Set 2 (8A10).

Variable format for item pool, using I-fields. It must be contained on one card (see Item Pool below).

Card Set 3 (16I5).

Columns 1-5: The number of items to be omitted, i.e., excluded from the computations. If none, punch "0" in column 5.  
Columns 6-10 and subsequent five-column fields: The item identification numbers of items to be omitted. Continue on as many cards as necessary. If more than one card is necessary, begin punching on the second card in columns 1-5.

Card Set 4 (8A10).

Variable input format for reading subject information and test response data. It must be contained on one card (see Test Response Data below for field type specifications).

Card Set 5 (8A10).

Description of the run is written on three cards. These cards are required, even if they are left blank.

LINPSCO (Card Set 2-7).

Card Set 2 (8A10).

Variable format for the item pool. It must be contained on one card (see Item Pool below for field type specifications).

Card Set 3 (80I1).

Punch in the  $n^{\text{th}}$  column the number of response categories minus 1 for the  $n^{\text{th}}$  item. Continue on as many cards as necessary.

Card Set 4 (16I5).

Punch in five-column fields the item identification numbers of the items in the test. The numbers must appear in the same order as the items appear in the test. Continue on as many cards as necessary.

Card Set 5 (80I1).

The information on this card is used to omit specified items from the computations. To omit the  $n^{\text{th}}$  item, punch a "1" in the  $n^{\text{th}}$  column of this card; otherwise, punch "0." If no items are to be omitted, punch as many zeros as there are items in the test. Continue on as many cards as necessary.

Card Set 6 (8A10).

Variable format for subject information and test response data. It must be contained on one card (see Test Response Data below for field type specifications).

Card Set 7 (8A10).

Description of the run is written on three cards.

### Item Pool

LINDSCO and ADADSCO. To score the response data, a file containing the item pool item parameter estimates must be prepared beforehand and placed in a file called IPOOL. The file consists of a line for each item in the pool with the following information:

1. A unique item number;
2. Estimate of the  $a$  parameter;
3. Estimate of the  $b$  parameter;
4. Estimate of the  $c$  parameter;
5. Correct alternative for this item, i.e., the keyed response.

For LINDSCO, only Items 1 through 4 must be supplied; for ADADSCO, Item 5 must be supplied also, although it could be a "dummy" key (e.g., a blank), since the data may already be scored (see columns 44-45 for Card Set 1).

The exact format of this information is not critical, since it is read with a user-specified variable format. However, the following limitations must be observed: (1) the information must be read in the above order; (2) the item number must be read in integer mode; (3) the item parameter estimates must be read in floating point; and (4) the key, if ADADSCO is being used, must be read in integer mode.

A typical format for LINDSCO could be  
(10X,I4,3F10.2) .

For ADADSCO, a typical format might be  
(10X,I4,3F10.2,I2) .

All three parameter estimates must be read even if the user is using a one- or two-parameter model. This presents no difficulties, however, since in the case of, say, a two-parameter model, the third parameter is 0 for all items. This may be accomplished by reading blanks or zeros, or by editing item parameter estimates (see below).

The number of items in the pool may range from the number of items in the test,  $M$ , to 600. If the item pool for LINDSCO consists of only the items being scored in the test, then OPT2 should be set to 1. This indicates to the program that items do not have to be searched. On the other hand, if the pool consists of items in addition to those used in the present test, then OPT2 should be set to 0. This instructs the program to search for the item and to retrieve the corresponding item parameters. For both LINDSCO and ADADSCO, if at least one of the items being called for is not found in the pool, the program prints a message; and the unavailable item is treated as an omitted item.

Editing of item parameter estimates. LINDSCO and ADADSCO have several options to edit item parameter estimates. If OPT3=1, the program checks that the item parameter estimates are within certain bounds. For the discrimination ( $a$ ) parameter, the program checks to see if the estimate exceeds AMAX; if it does, it is set to AMAX. For the difficulty ( $b$ ) parameter, if the estimate is below BMIN, it is set to BMIN; if it is above BMAY, it is set to BMAX. For the "guessing" ( $c$ ) parameter, the program checks to see if the estimate exceeds CMAX; if it does, it is set to CMAX. If the user wants to edit only one or two parameters, the limits of the other parameters should be chosen so that the editing has no effect.



A more radical form of editing is also possible. If OPT3=2, then in addition to the editing caused by OPT3=1, the program sets all  $c$  parameter estimates to CMAX. If CMAX=0.0, this implies that a two-parameter model is in effect. If OPT3=3, then in addition to the editing caused by OPT3=1 and OPT3=2, the program sets all  $a$  parameter estimates to AMAX.

LINPSCO. For polychotomous scoring, the item pool consists of the following information for graded normal and logistic ogive models:

1. A *unique* item identification number;
2. The "discrimination" parameter, which is common to all response categories;
3.  $m_g - 1$  "difficulty" parameters, where  $m_g$  is the number of response categories in the  $g^{\text{th}}$  item. Since  $m_g$  can be at most 10, there would be at most 9 difficulty parameters.

The exact format for reading this information is not crucial, since it is read by a user-supplied format statement. However, the following restrictions must be observed: (1) the identification number is read first, in integer mode; (2) next, the estimated discrimination parameter is read in floating point mode; (3) the  $m_g - 1$  "difficulty" parameters are read next, with the difficulty of the best alternative followed by the second best alternative, and so forth.

Since the program allows the number of categories to differ from item to item, the format should be specified so that it can read the information for the item with the most response categories. For example, if in a given test, the maximum number of response categories is seven, then there should be at most six difficulty parameters. The format for such pools might be as follows:

(I4,6X,F5.2/10X,6F5.2) .

In this format the item identification number is read in the I4 field; the discrimination parameter is read next in format F5.2; and the six difficulty values are read from the next card, beginning in column 11.

For the nominal logistic model, the item information is read in the following order:

1. A *unique* item identification number;
2.  $m_g$  "slope" parameters; and
3.  $m_g$  intercept parameters.

Differing from the graded models, in the nominal model there is a pair of parameters (a slope and an intercept) associated with *each* response category. Since the response categories are not ordered in the nominal model, the order in which the parameters are read is unimportant. However, the ordinal position in which the parameters appear in the pool must correspond with the integer associated with that response category. As in the graded models, the format should be able to read the information for the item with most response categories. For example, if the maximum number of response categories is five, the format

could be

(I4,16X,5F5.2,5X,5F5.2) .

In this format, the item identification number is read in the I4 field; next, the five slope parameters are read in 5F5.2; and finally, the five intercept parameters are read in the last set of 5F5.2 fields.

#### Test Response Data

Data for all testees must be on a file called DATA. The structure of this file differs slightly for each of the programs. In all cases, however, the last record of DATA must be an end-of-record marker.

LINDSCO. This program requires that for each individual the following information be provided on DATA:

1. Name,
2. Identification number, and
3. Responses to the test items.

The exact format of this information is not critical, since it is read with a user-supplied variable format; but the information must appear in the above order. Two words are used for testee name; thus, name should be read with two alphanumeric words, e.g., 2A10. This allows for up to 20 characters. The testee identification is read with an alphanumeric field of at least 1 column, e.g., A1, A9. Test item responses are read with an integer format, e.g., 2011.

The test data may be raw item responses (i.e., the number of the alternatives chosen) or scored (i.e., 0 for incorrect and 1 for correct). However, in either case, a scoring key must be provided (see Card Set 5 for LINDSCO). The key will contain the number of the correct alternative if raw data are read. If the data are already scored, a "dummy" key full of "1's" must be provided.

Omitted items are indicated by the integer IOMIT (see columns 46 and 47 of Card Set 1 for LINDSCO). For raw data, this will normally be an integer greater than the number of alternatives. Similarly, for scored (0-1) data IOMIT must be an integer greater than 1.

ADADSCO. The program requires that the following information be provided on DATA for each individual:

1. Name,
2. Identification number,
3. Number of items answered by the testee (i.e., number of items attempted),
4. Item identification numbers of items attempted, and
5. Responses to the test items.

This information is read in the above order with a user-supplied variable format; thus, the exact format is not critical. However, the following limitations must be observed. Even though the number of items administered usually varies across individuals in an adaptive test, this program assumes that the data record for each testee is formally the same (i.e., that there is the same number of data

lines per testee and that these lines contain similar information). Thus, if the maximum number of items taken by anyone is MMAX (see Card Set 1 for ADADSCO), but any particular testee takes  $M$  items, where  $M < \text{MMAX}$ , then that testee's record should be "padded" to MMAX items. This can be accomplished by leaving an appropriate number of blank fields. The name is dimensioned for two words so the format should allow for two words, e.g., 2A10. The identification number is read with an alphanumeric format, e.g., A8. The number of items is read in integer mode. The item identification numbers and item responses are also read in integer mode. Note that in reading the item identification numbers and the item responses, the format should read MMAX of each, even if some of these will be blank for a given individual.

As an example assume that MMAX was 25; then the variable format could be (2A10,A10,I2/20I4/5I4/25I1).

In this format, the name, testee identification, and  $M$  are read from the first card; the item identification numbers are read from the next two cards; and finally, the item responses are read from the fourth card. Note that for testees attempting 20 items or less, the third card will be blank.

The item responses may be scored or raw data. For scored data, the responses have been reduced to three categories: correct, incorrect, and omitted. In this case, IFLAG should be set to the integer corresponding to the correct code, and IOMIT should be set to the code for omitted responses. Note that if  $\text{IFLAG} > 0$ , the program ignores the key read as part of the item pool. For raw data, the key will have been read as part of the item pool; IFLAG must therefore be set to 0. IOMIT will still be operational, however; and it must be set to an integer other than the highest numbered response alternative.

LINPSCO. The DATA file is similar to LINDSCO's with the exception that the item responses must include the response category chosen by the testee for a given item. For graded models, the convention that the best response category be coded "1," second best "2," and so forth, must be obeyed. For the nominal logistic model, this convention does not apply; but care must be taken so that a category's response code matches the ordinal position of that category in the IPOOL file. For either graded or nominal data, the code for omitted responses should be an integer greater than the maximum number of response categories.

### Output

Four kinds of output are produced by each program: program parameters, item parameters, computational messages, and testee data.

#### Program Parameters

The output consists of the information in Card Set 1, the description of the run, and the variable formats for reading the item pool and the testee's raw data.

#### Item Parameters

LINDSCO and LINPSCO. The output consists of item identification number, scoring key, rejection key (i.e., whether or not the item was included in the computations), and the item parameter estimates. If the estimates have been



edited, the edited values will be printed. An option (see column 10 of Card Set 1) permits all of this information to be punched as well. If subscale scoring has been requested, the item identification number of the items in each subscale will be printed.

ADADSCO. The user has the option, but only for the first 10 testees, to print the following: testee's name and identification number; and for each item attempted, the item identification number, the response to that item, and the item parameter estimates.

#### Computational Messages

The program will print a testee's name and identification number if (1) a response pattern is found with all items correct or incorrect, excluding omitted or rejected items; (2) a zero score has been obtained; or (3) it was not possible to achieve convergence in scoring the testee's responses. For polychotomous data, a perfect or zero vector occurs if the testee responds with the best or worst response categories in all attempted items, exclusive of omitted or rejected items. If an item is not found in the pool or has extreme parameter estimates, an informative message is printed.

The number of testees read and the number of convergence failures are also printed. If Bayesian scoring has been requested, the number of nonconvergent cases will be zero.

#### Testee Data

LINDSCO. For each testee, the following information is written on a file called TAPE3:

1. Name;
2. Testee identification number;
3. Scale number, or in the case of total score, a "T";
4. Proportion of items answered correctly;
5. Maximum likelihood or Bayesian estimate of  $\theta$ ;
6. The response pattern information for maximum likelihood scoring or the posterior variance of  $\theta$  for Bayesian scoring;
7. The number of items used in the estimation of  $\theta$ , excluding items rejected, omitted, or not found;
8. The test information associated with the estimated  $\theta$  (for Bayesian scoring, the information is computed using the normal ogive model);
9. The true score corresponding to the estimated  $\theta$ ;
10. For maximum likelihood scoring,
  - a. The number of Newton-Raphson iterations needed to achieve convergence and
  - b. The standard error of  $\theta$ .

The format used for writing this information for total scores is  
(X,2A10,A9,\*T\*,F5.2,2F7.2,I4,2F7.2,I4,F7.2) .

The subscale results are written with  
(X,2A10,A9,I2,F5.2,2F7.2,I4,2F7.2,I4,F7.2) .

ADADSCO. The same information is written as that for LINDSCO with the exception of the scale number. The format is  
(X,2A10,A9,F5.2,2F7.2,I4,2F7.2,I4,F7.2) .

LINPSCO. For LINPSCO, the following information is written:

1. Name;
2. Testee identification number;
3. Proportion of "best" responses;
4. Maximum likelihood estimate of  $\theta$ ;
5. The response pattern information;
6. The number of items used in the estimation of  $\theta$  excluding items rejected, omitted, or not found;
7. The number of iterations needed to achieve convergence.
8. The test information associated with the estimated  $\theta$ ;
9. Estimated standard error of measurement.

#### AVAILABILITY

FORTRAN source code listings of the three programs are in Appendix C (LINDSCO), Appendix D (ADADSCO), and Appendix E (LINPSCO). Copies of the FORTRAN source code are available on cards or tape at nominal cost from

Psychometric Methods Program  
Department of Psychology  
University of Minnesota  
75 East River Road  
Minneapolis, Minnesota 55455

Telephone: 612-376-7378

Potential users of these programs should note that the programs were written for Control Data Corporation CYBER series computers. Because of the large word size of the CYBER computers, accurate computation on other computers may require the use of double-precision arithmetic. Minimal additional modifications required may include (1) modification of A10 fields to smaller sizes used by other computers and (2) modification of FORTRAN statements unique to the CYBER series computers.

#### REFERENCES

- Bejar, I. I., Weiss, D. J., & Gialluca, K. A. An information comparison of conventional and adaptive tests in the measurement of classroom achievement (Research Report 77-7). Minneapolis: University of Minnesota, Department of Psychology, Psychometric Methods Program, October 1977. (NTIS No. AD A047495)
- Birnbaum, A. Some latent trait models and their use in inferring an examinee's ability. In F. M. Lord & M. R. Novick, Statistical theories of mental test scores. Reading, MA: Addison-Wesley, 1968.
- Bock, R. D. Estimating latent parameter and ability when responses are scored on two or more nominal categories. Psychometrika, 1972, 37, 29-51.
- Kendall, M. G., & Stuart, A. The advanced theory of statistics (Vol 2). New York: Hafner, 1961.
- Kolakowski, D., & Bock, R. D. A FORTRAN IV program for maximum likelihood item analysis and test scoring: Normal ogive model (Research Memo No. 12). Chicago: University of Chicago, Department of Education, Statistics Laboratory, 1970.
- Kolakowski, D., & Bock, R. D. LOGOG: Maximum likelihood item analysis and test scoring: Logistic model for multiple responses (Research Memo No. 13). Chicago: University of Chicago, Department of Education, Statistics Laboratory, 1972.
- Lawley, D. N. On problems connected with item selection and test construction. Proceedings of the Royal Society of Edinburgh, 1943, 61, 273-287.
- Lord, F. M. Some test theory for tailored testing. In W. H. Holtzman (Ed.), Computer-assisted instruction, testing, and guidance. New York: Harper & Row, 1970.
- Lord, F. M., & Novick, M. R. Statistical theories of mental test scores. Reading, MA: Addison-Wesley, 1968.
- Owen, R. J. A Bayesian sequential procedure for sequential response in the context of adaptive mental testing. Journal of the American Statistical Association, 1975, 70, 351-356.
- Samejima, F. Estimation of latent ability using a response pattern of graded scores. Psychometrika, Monograph Supplement No. 17, 1969.
- Samejima, F. A comment on Birnbaum's three-parameter logistic model in the latent trait theory. Psychometrika, 1973, 28, 221-234.
- Sympson, J. B. Estimation of latent trait status in adaptive testing procedures. In D. J. Weiss (Ed.), Applications of computerized adaptive testing (Research Report 77-1). Minneapolis: University of Minnesota, Department of Psychology, Psychometric Methods Program, March 1977. (NTIS No. AD A038114)



Weiss, D. J. (Ed.) Computerized adaptive trait measurement: Problems and prospects (Research Report 75-5). Minneapolis: University of Minnesota, Department of Psychology, Psychometric Methods Program, November 1975. (NTIS No. AD A018675)

Weiss, D. J. Computerized ability testing, 1972-1975 (Final Report). Minneapolis: University of Minnesota, Department of Psychology, Psychometric Methods Program, April 1976. (NTIS No. A024516).

Wood, R. L., Wingersky, M. S., & Lord, F. M. LOGIST: A computer program for estimating examinee ability and item characteristic curve parameters (Research Memo 76-6). Princeton, NJ: Educational Testing Service, 1976.

Wright, B. D., & Mean, R. J. CALFIT: Sample free item calibration with a Rasch measurement model (Research Memo No. 18). Chicago: University of Chicago, Department of Education, Statistical Laboratory, 1976.

# APPENDICES

## Appendix A

### Item Parameter Estimation Programs

Program Name	Model	Reference
LOGIST	Three-parameter logistic ogive	Wood, R. L., Wingersky, M. S., & Lord, F. M. LOGIST: A computer program for estimating examinee ability and item characteristic curve parameters (Research Memorandum 76-6). Princeton, NJ: Educational Testing Service, 1976.
NORMOG	Three-parameter normal ogive	Kolakowski, D., Bock, R. A. A FORTRAN IV program for maximum likelihood item analysis and test scoring: Normal ogive model (Research Memo No. 12). Chicago: University of Chicago, Department of Education, Statistics Laboratory, 1970.
ESTEM	Three-parameter logistic or normal ogive	No program documentation available. For a description of the procedure see Urry, V. W., Ancillary estimators for the parameters of mental test models. Washington, DC: U.S. Civil Service Commission, Personnel Research and Development Center, 1974.
BICAL	One-parameter logistic ogive	Wright, B. D., & Mead, R. J. CALFIT: Sample-free item calibration with a Rasch measurement model (Research Memorandum No. 18). Chicago: University of Chicago, Department of Education, Statistical Laboratory, 1976.
LOGOG	Graded normal and logistic ogive, nominal logistic	Kolakowski, D., & Bock, R. D. LOGOG: A FORTRAN IV program for maximum likelihood item analysis and test scoring: Logistic model for multiple response (Research Memorandum No. 13). Chicago: University of Chicago, Department of Education, Statistical Laboratory, 1972.

*Appendix B:*  
*Examples of Program Use*

The following examples serve to illustrate the use of each of the three programs. These results should also be useful in testing the accuracy of the results of the programs in different installations.

LINDSCO

The IPOOL file for these examples was

1	1.000	-2.000	.25
2	1.500	-1.000	.25
3	1.000	0.000	.25
4	1.500	1.000	.25
5	1.000	2.000	.25

The DATA file is also shown below. The first field contains the names; the second, the subject identification; and the third, the response patterns.

Name	I.D.	Responses
A00	1	00000
A01	2	00001
A02	3	00010
A03	4	00011
A04	5	00100
A05	6	00101
A06	7	00110
A07	8	00111
A08	9	01000
A09	10	01001
A10	11	01010
A11	12	01011
A12	13	01100
A13	14	01101
A14	15	01110
A15	16	01111
A16	17	10000
A17	18	10001
A18	19	10010
A19	20	10011
A20	21	10100
A21	22	10101
A22	23	10110
A23	24	10111
A24	25	11000
A25	26	11001
A26	27	11010
A27	28	11011
A28	29	11100
A29	30	11101
A30	31	11110
A31	32	11111



Example 1. This example illustrates the use of the normal ogive model (OPT4=1) for a five-item test ( $m=5$ ) with a pool containing five items (INUP=5). The example also illustrates the use of parameter editing (OPT3=1) in which BMAX=BMIN=0.0, which in effect sets all  $b$  parameter estimates to 0.0. AMAX=2.00, which means that if there were  $a$  parameter estimates greater than 2, they would be set to 2.00. CMAX=.10, which means that any  $c$  parameter estimates greater than .10 will be edited to .10. This example also illustrates the use of subscales. The program parameter cards for this example were

```

      5      5      111 0.00 1.00 2.00 0.00 0.00  .10  4444
(9X,I1,3F10.3)
      1      2      3      4      5
00000
11111
(2A4,A2,10X,5I1)
RUNS BASED ON ALL POSSIBLE RESPONSE VECTORS FOR A FIVE ITEM TEST

```

```

      1
      3      1
      2      3      5

```

The output corresponding to this example is shown on the following pages.

LINDSCO

LINEAR DICHOTOMUS SCORING WITH THREE PARAMETER MODELS

PSYCHOMETRIC METHODS PROGRAM  
DEPARTMENT OF PSYCHOLOGY  
UNIVERSITY OF MINNESOTA  
MPLS. MINN. 55455

INUP = 5  
MMAX = 5  
IOMI1 = 4444  
OPT1 = -0  
OPT2 = 1  
OPT3 = 1  
OPT4 = 1  
TS = 0  
TSS = 1.00  
AMAX = 2.00  
BMAX = 0  
BMIN = 0  
CMAX = .10  
VARIABLE FORMAT FOR POOL=(9X,I1,3F10.3)  
VARIABLE FORMAT FOR DATA=(2A4,A2,10X,5I1)  
RUNS BASED ON ALL POSSIBLE RESPONSE VECTORS FOR A FIVE ITEM TEST

ITEMS IN SUBSCALE NO= 1

2 3 5

ITEM	ID	S	KEYS	REJECTIONS	A	B	C
1	1	0	1.00	0	.10		
2	1	0	1.50	0	.10		
3	1	0	1.00	0	.10		
4	1	0	1.50	0	.10		
5	1	0	1.00	0	.10		

SUBJECT =A00	ID = 1	HAS NO ANSWERS CORRECT IN TOTAL SCALE	1
SUBJECT =A00	ID = 3	HAS NO ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A02	ID = 14	HAS ALL ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A13	ID = 16	HAS ALL ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A15	ID = 17	HAS NO ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A16	ID = 19	HAS NO ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A18	ID = 30	HAS ALL ANSWERS CORRECT IN SUBSCALE	1
SUBJECT =A29	ID = 32	HAS ALL ANSWERS CORRECT IN TOTAL SCALE	1
SUBJECT =A31	ID = 32	HAS ALL ANSWERS CORRECT IN SUBSCALE	1

CASES READ= 32 CASES NOT CONVERGED= 0

Contents of TAPE3:

Testee Name	Testee Identification Number	Subscale: T=Total; I=Scale 1	Proportion Correct	Maximum Likelihood Estimate of $\theta$	Response Pattern Information	Number of Items	Test Information Associated with $\hat{\theta}$	Expected Proportion Correct	Number of Iterations	Estimated Standard Error of Measurement
A00	1	T	0	-10.00	0	5	0	0	0	
A00	1	I	0	-10.00	0	5	0	0	0	
A01	2	T	.20	-1.06	1.48	5	1.07	.20	2	.82
A01	2	I	.33	-.65	1.51	3	1.36	.30	3	.81
A02	3	T	.20	-.95	.58	5	1.36	.22	5	1.31
A02	3	I	0	-10.00	0	5	0	0	0	
A03	4	T	.40	-.31	3.35	5	3.40	.42	3	.55
A03	4	I	.33	-.65	1.51	3	1.36	.30	3	.81
A04	5	T	.20	-1.06	1.48	5	1.07	.20	2	.82
A04	5	I	.33	-.65	1.51	3	1.36	.30	3	.81
A05	6	T	.40	-.50	3.08	5	2.84	.35	2	.57
A05	6	I	.67	.05	2.33	3	2.23	.57	2	.66
A06	7	T	.40	-.31	3.35	5	3.40	.42	3	.55
A06	7	I	.33	-.65	1.51	3	1.36	.30	3	.81
A07	8	T	.60	.07	3.98	5	3.94	.58	2	.50
A07	8	I	.67	.05	2.33	3	2.23	.57	2	.66
A08	9	T	.20	-.95	.58	5	1.36	.22	5	1.31
A08	9	I	.33	-.30	1.83	3	1.95	.43	3	.74
A09	10	T	.40	-.31	3.35	5	3.40	.42	3	.55
A09	10	I	.67	.40	2.03	3	2.11	.71	2	.70
A10	11	T	.40	-.12	3.63	5	3.78	.50	2	.52
A10	11	I	.33	-.30	1.83	3	1.95	.43	3	.74
A11	12	T	.60	.27	3.71	5	3.88	.67	2	.52
A11	12	I	.67	.40	2.03	3	2.11	.71	2	.70
A12	13	T	.40	-.31	3.35	5	3.40	.42	3	.55
A12	13	I	.67	.40	2.03	3	2.11	.71	2	.70
A13	14	T	.60	.07	3.98	5	3.94	.58	2	.50
A13	14	I	1.00	10.00	0	5	0	0	0	
A14	15	T	.60	.27	3.71	5	3.88	.67	2	.52
A14	15	I	.67	.40	2.03	3	2.11	.71	2	.70
A15	16	T	.80	.75	2.82	5	2.94	.83	2	.60
A15	16	I	1.00	10.00	0	5	0	0	0	
A16	17	T	.20	-1.06	1.48	5	1.07	.20	2	.82
A16	17	I	0	-10.00	0	5	0	0	0	
A17	18	T	.40	-.50	3.08	5	2.84	.35	2	.57
A17	18	I	.33	-.65	1.51	3	1.36	.30	3	.81
A18	19	T	.40	-.31	3.35	5	3.40	.42	3	.55
A18	19	I	0	-10.00	0	5	0	0	0	
A19	20	T	.60	.07	3.98	5	3.94	.58	2	.50
A19	20	I	.33	-.65	1.51	3	1.36	.30	3	.81
A20	21	T	.40	-.50	3.08	5	2.84	.35	2	.57
A20	21	I	.33	-.65	1.51	3	1.36	.30	3	.81
A21	22	T	.60	-.11	3.94	5	3.79	.50	2	.50
A21	22	I	.67	.05	2.33	3	2.23	.57	2	.66
A22	23	T	.60	.07	3.98	5	3.94	.58	2	.50
A22	23	I	.33	-.65	1.51	3	1.36	.30	3	.81
A23	24	T	.80	.46	3.82	5	3.62	.74	2	.51
A23	24	I	.67	.05	2.33	3	2.23	.57	2	.66
A24	25	T	.40	-.31	3.35	5	3.40	.42	3	.55
A24	25	I	.33	-.30	1.83	3	1.95	.43	3	.74
A25	26	T	.60	.07	3.98	5	3.94	.58	2	.50
A25	26	I	.67	.40	2.03	3	2.11	.71	2	.70
A26	27	T	.60	.27	3.71	5	3.88	.67	2	.52
A26	27	I	.33	-.30	1.83	3	1.95	.43	3	.74
A27	28	T	.80	.75	2.82	5	2.94	.83	2	.60
A27	28	I	.67	.40	2.03	3	2.11	.71	2	.70
A28	29	T	.60	.07	3.98	5	3.94	.58	2	.50
A28	29	I	.67	.40	2.03	3	2.11	.71	2	.70
A29	30	T	.80	.46	3.82	5	3.62	.74	2	.51
A29	30	I	1.00	10.00	0	5	0	0	0	
A30	31	T	.80	.75	2.82	5	2.94	.83	2	.60
A30	31	I	.67	.40	2.03	3	2.11	.71	2	.70
A31	32	T	1.00	10.00	0	5	0	0	0	
A31	32	I	1.00	10.00	0	5	0	0	0	



Example 2. This example is identical to Example 1 except that the Bayesian scoring routine was used (OPT4=3) instead of the maximum likelihood normal ogive. Only the scoring results are shown.

Testee Name	Testee Identification Number	Subscale: T=Total; I=Scale 1	Proportion correct	Bayesian estimate of $\theta$	Bayesian Posterior Variance	Number of Items	Test Information Associated with $\theta$	Expected Proportion Correct			
AU0	1	T	0	-1.23	.31	5	.71	.17	0	0	0
AU0	1	I	0	-1.04	.39	3	.69	.21	0	0	0
AU1	2	T	.20	-.88	.35	5	1.58	.24	0	0	0
AU1	2	I	.33	-.55	.44	3	1.55	.34	0	0	0
AU2	3	T	.20	-.83	.38	5	1.73	.25	0	0	0
AU2	3	I	0	-1.04	.39	3	.69	.21	0	0	0
AU3	4	T	.40	-.34	.41	5	3.33	.41	0	0	0
AU3	4	I	.33	-.55	.44	3	1.55	.34	0	0	0
AU4	5	T	.20	-.89	.30	5	1.56	.23	0	0	0
AU4	5	I	.33	-.55	.41	3	1.54	.33	0	0	0
AU5	6	T	.40	-.51	.32	5	2.83	.35	0	0	0
AU5	6	I	.67	.02	.44	3	2.22	.56	0	0	0
AU6	7	T	.40	-.37	.33	5	3.23	.40	0	0	0
AU6	7	I	.33	-.55	.41	3	1.54	.33	0	0	0
AU7	8	T	.60	.11	.35	5	3.95	.60	0	0	0
AU7	8	I	.67	.02	.44	3	2.22	.56	0	0	0
AU8	9	T	.20	-.76	.28	5	1.92	.26	0	0	0
AU8	9	I	.33	-.55	.39	3	1.87	.41	0	0	0
AU9	10	T	.40	-.40	.30	5	3.14	.38	0	0	0
AU9	10	I	.67	.23	.43	3	2.22	.65	0	0	0
A10	11	T	.40	-.25	.30	5	3.53	.44	0	0	0
A10	11	I	.33	-.55	.39	3	1.87	.41	0	0	0
A11	12	T	.60	.20	.33	5	3.93	.64	0	0	0
A11	12	I	.67	.23	.43	3	2.22	.65	0	0	0
A12	13	T	.40	-.40	.27	5	3.15	.38	0	0	0
A12	13	I	.67	.22	.42	3	2.22	.64	0	0	0
A13	14	T	.60	-.02	.28	5	3.89	.54	0	0	0
A13	14	I	1.00	.98	.50	3	1.35	.88	0	0	0
A14	15	T	.60	.14	.30	5	3.93	.63	0	0	0
A14	15	I	.67	.22	.42	3	2.22	.64	0	0	0
A15	16	T	.60	.69	.33	5	3.10	.81	0	0	0
A15	16	I	1.00	.96	.50	3	1.35	.88	0	0	0
A16	17	T	.20	-.82	.25	5	1.76	.25	0	0	0
A16	17	I	0	-1.04	.39	3	.69	.21	0	0	0
A17	18	T	.40	-.50	.26	5	2.84	.35	0	0	0
A17	18	I	.33	-.55	.44	3	1.55	.34	0	0	0
A18	19	T	.40	-.31	.27	5	3.19	.39	0	0	0
A18	19	I	0	-1.04	.39	3	.69	.21	0	0	0
A19	20	T	.60	.01	.29	5	3.91	.55	0	0	0
A19	20	I	.33	-.55	.44	3	1.55	.34	0	0	0
A20	21	T	.40	-.50	.24	5	2.84	.35	0	0	0
A20	21	I	.33	-.55	.41	3	1.54	.33	0	0	0
A21	22	T	.60	-.17	.25	5	3.69	.48	0	0	0
A21	22	I	.67	.02	.44	3	2.22	.56	0	0	0
A22	23	T	.60	-.01	.26	5	3.90	.55	0	0	0
A22	23	I	.33	-.55	.41	3	1.54	.33	0	0	0
A23	24	T	.60	.39	.27	5	3.73	.71	0	0	0
A23	24	I	.67	.02	.44	3	2.22	.56	0	0	0
A24	25	T	.40	-.37	.24	5	3.23	.40	0	0	0
A24	25	I	.33	-.55	.39	3	1.87	.41	0	0	0
A25	26	T	.60	-.02	.25	5	3.89	.54	0	0	0
A25	26	I	.67	.23	.43	3	2.22	.65	0	0	0
A26	27	T	.60	.17	.27	5	3.94	.62	0	0	0
A26	27	I	.33	-.55	.39	3	1.87	.41	0	0	0
A27	28	T	.60	.61	.30	5	3.29	.79	0	0	0
A27	28	I	.67	.23	.43	3	2.22	.65	0	0	0
A28	29	T	.60	-.03	.24	5	3.88	.54	0	0	0
A28	29	I	.67	.22	.42	3	2.22	.64	0	0	0
A29	30	T	.60	.34	.26	5	3.80	.69	0	0	0
A29	30	I	1.00	.96	.50	3	1.35	.88	0	0	0
A30	31	T	.60	.59	.32	5	3.33	.78	0	0	0
A30	31	I	.67	.22	.42	3	2.22	.64	0	0	0
A31	32	T	1.00	1.20	.37	5	1.72	.92	0	0	0
A31	32	I	1.00	.98	.50	3	1.35	.88	0	0	0

Example 3. This example illustrates the use of the maximum likelihood logistic scoring routine (e.g., OPT4=2) without subscale scoring. Only the scoring results are shown.

Testee Name	Testee Identification Number	Subscale: T=Total; I=Subscale 1	Proportion Correct	Maximum Likelihood Estimate of $\theta$	Response Pattern Information	Number of Items	Test Information Associated with $\theta$	Expected Proportion Correct	Number of Iterations	Estimated Standard Error of Measurement
AU0	1	T	0	-10.00	0	5	0	0	0	
AU1	2	T	.20	-1.00	1.20	5	.90	.20	2	.91
AU2	3	T	.20	-99.99	-99.99	5	-99.99	-99.99	99	-99.99
AU3	4	T	.40	-.29	3.59	5	3.68	.42	3	.53
AU4	5	T	.20	-1.05	1.20	5	.90	.20	2	.91
AU5	6	T	.40	-.46	3.32	5	2.95	.35	1	.55
AU6	7	T	.40	-.29	3.59	5	3.68	.42	3	.53
AU7	8	T	.60	.07	4.49	5	4.46	.58	2	.47
AU8	9	T	.20	-99.99	-99.99	5	-99.99	-99.99	99	-99.99
AU9	10	T	.40	-.29	3.59	5	3.68	.42	3	.53
A10	11	T	.40	-.11	4.01	5	4.26	.50	2	.50
A11	12	T	.60	.25	4.22	5	4.26	.66	2	.49
A12	13	T	.40	-.29	3.59	5	3.68	.42	3	.53
A13	14	T	.60	.07	4.49	5	4.46	.58	2	.47
A14	15	T	.60	.25	4.22	5	4.26	.66	2	.49
A15	16	T	.80	.71	2.74	5	2.72	.83	2	.60
A16	17	T	.20	-1.05	1.20	5	.90	.20	2	.91
A17	18	T	.40	-.46	3.32	5	2.95	.35	1	.55
A18	19	T	.40	-.29	3.59	5	3.68	.42	3	.53
A19	20	T	.60	.07	4.49	5	4.46	.58	2	.47
A20	21	T	.40	-.46	3.32	5	2.95	.35	1	.55
A21	22	T	.60	-.10	4.53	5	4.28	.50	2	.47
A22	23	T	.60	.07	4.49	5	4.46	.58	2	.47
A23	24	T	.80	.45	3.70	5	3.70	.74	2	.52
A24	25	T	.40	-.29	3.59	5	3.68	.42	3	.53
A25	26	T	.60	.07	4.49	5	4.46	.58	2	.47
A26	27	T	.60	.25	4.22	5	4.26	.66	2	.49
A27	28	T	.80	.71	2.74	5	2.72	.83	2	.60
A28	29	T	.60	.07	4.49	5	4.46	.58	2	.47
A29	30	T	.80	.45	3.70	5	3.70	.74	2	.52
A30	31	T	.80	.71	2.74	5	2.72	.83	2	.60
A31	32	T	1.00	10.00	0	5	0	0	0	

ADADSCO

IPOOL for this example consisted of 10 items:

1	1.00	-2.00	.25 1
2	1.25	-1.50	.25 1
3	1.50	-1.00	.25 1
4	1.75	-0.50	.25 1
5	1.00	0.00	.25 1
6	1.25	0.50	.25 1
7	1.50	1.00	.25 1
8	1.75	1.50	.25 1
9	1.00	2.00	.25 1
10	1.25	2.50	.25 1

The data for the 16 subjects used in the example are shown below:

```

1011101110
1 2 3 4 5 6 7 8 9 10
2 B 4
1010
2 4 6 8
3 C 5
11111
1 2 3 4 5
4 D 5
00000
6 7 8 9 10
5 E 8
11000000
1 2 3 5 6 7 8 9
6 F 2
10
3 7
7 G 6
011111
1 2 4 6 8 9
8 H 9
101010101
1 2 3 4 5 6 7 8 9
9 I 7
1100110
1 3 4 5 6 8 10
10 J 3
110
4 8 9

```



```
11 K      8
10110110
 2 3 4 6 7 8 910
12 L      9
101111111
 1 2 3 4 5 7 8 910
13 M      5
01001
 1 3 5 7 9
14 N      6
000001
 4 5 6 7 8 9
15 P      7
1011110
 1 2 4 5 6 7 8
16 Q      6
011011
 2 4 5 7 810
```

The program control cards for this example were

```
10 10 1012 0.00 1.00 2.00 0.00 0.00 .10 1 3
(8X,12,3F10.2,12)
0
(A2,1X,2A2,12,/10I1,/10I2)
DESCRIPTION
```

In this example the maximum number of items attempted by anyone was 10 (MMAX=10). Although the code for omitted items was 3 (IOMIT=3), IFLAG=1, which means the key to each item was read from IPOOL; however, in this case it was 1 for all items. OPT1=1 means that item information for the first 10 subjects will be printed. Editing of item parameters was requested (OPT3=1). The scoring algorithm was maximum likelihood logistic.

The entire output for this example is shown on the following pages.

ADADSCG  
=====

ADAPTIVE DICHOTOMOUS SCORING WITH THREE PARAMETER MODELS

PSYCHOMETRIC METHODS PROGRAM  
DEPARTMENT OF PSYCHOLOGY  
UNIVERSITY OF MINNESOTA  
MPLS. MINN. 55455

INUP = 10  
NMAX = 10  
IOMIT = 3  
IFLAG = 1  
OPT1 = 1  
OPT2 = 0  
OPT3 = 1  
OPT4 = 2  
IS = 0  
TSS = 1.00  
AMAX = 2.00  
BMAX = 0  
BMIN = 0  
CMAX = .10  
VARIABLE FORMAT FOR POOL=(8X,I2,3F10.2,I2)  
VARIABLE FORMAT FOR DATA=(A2,1X,2A2,I2,/,10I1,/,10I2)  
DESCRIPTION

Item Identification Number	Scored Answer	Testee Number	Discrimination Parameter (a)	Difficulty Parameter (b)	"Guessing" Parameter (c)
1	1	1	1.00	0	.10
2	0	1	1.25	0	.10
3	1	1	1.50	0	.10
4	1	1	1.75	0	.10
5	1	1	1.00	0	.10
6	0	1	1.25	0	.10
7	1	1	1.50	0	.10
8	1	1	1.75	0	.10
9	1	1	1.00	0	.10
10	0	1	1.25	0	.10
2	1	2	1.25	0	.10
4	0	2	1.75	0	.10
6	1	2	1.25	0	.10
8	0	2	1.75	0	.10
1	1	3	1.00	0	.10
2	1	3	1.25	0	.10
3	1	3	1.50	0	.10
4	1	3	1.75	0	.10
5	1	3	1.00	0	.10
SUBJECT C					
6	0	4	1.25	0	.10
7	0	4	1.50	0	.10
8	0	4	1.75	0	.10
9	0	4	1.00	0	.10
10	0	4	1.25	0	.10

ID= 3 HAS ALL ANSWERS RIGHT

SUBJECT D  
E

IO= 4 HAS NO RIGHT ANSWERS

1	1	1.00	0	.10
2	1	1.25	0	.10
3	0	1.50	0	.10
5	0	1.00	0	.10
6	0	1.25	0	.10
7	0	1.50	0	.10
8	0	1.75	0	.10
9	0	1.00	0	.10
F				
3	1	1.50	0	.10
7	0	1.50	0	.10
G				
1	0	1.00	0	.10
2	1	1.25	0	.10
4	1	1.75	0	.10
6	1	1.25	0	.10
8	1	1.75	0	.10
9	1	1.00	0	.10
H				
1	1	1.00	0	.10
2	0	1.25	0	.10
3	1	1.50	0	.10
4	0	1.75	0	.10
5	1	1.00	0	.10
6	0	1.25	0	.10
7	1	1.50	0	.10
8	0	1.75	0	.10
9	1	1.00	0	.10
I				
1	1	1.00	0	.10
3	1	1.50	0	.10
4	0	1.75	0	.10
5	0	1.00	0	.10
6	1	1.25	0	.10
8	1	1.75	0	.10
10	0	1.25	0	.10
J				
4	1	1.75	0	.10
8	1	1.75	0	.10
9	0	1.00	0	.10

CASES READ= 16 CASES NOT CONVERGED= 0

Testee Name	Testee Identification Number	Proportion Correct	Maximum Likelihood Estimate of $\theta$	Response Pattern Information	Number of Items	Test Information Associated with $\theta$	Expected Proportion Correct	Number of Iterations	Estimated Standard Error of Measurement
A	1	.70	.34	9.65	10	9.65	.71	3	.32
B	2	.50	.22	5.06	4	4.66	.43	2	.44
C	3	1.00	10.00	0	5	0	0	0	0
D	4	0	-10.00	0	5	0	0	0	0
E	5	.25	-.78	3.15	8	2.69	.24	1	.56
F	6	.50	-.09	2.57	2	2.57	.50	2	.62
G	7	.83	.79	3.13	6	3.12	.87	2	.57
H	8	.56	-.09	9.85	9	9.55	.50	2	.32
I	9	.57	.05	8.03	7	8.03	.58	2	.35
J	10	.67	.43	3.18	3	3.20	.77	2	.56
K	11	.63	.13	9.62	8	9.62	.62	2	.32
L	12	.89	.89	4.02	9	4.02	.89	3	.50
M	13	.40	-.29	3.59	5	3.68	.42	3	.53
N	14	.17	-1.08	1.40	6	.85	.18	2	.84
P	15	.71	.25	7.55	7	7.52	.68	2	.36
Q	16	.67	.23	7.03	6	7.03	.67	2	.38



# LINPSCO

Following are sample runs from LINPSCO using graded models and the nominal logistic model.

Graded models. The IPOOL file for these examples was

1	1.5	3.0	2.0	4.5	5.0
2	1.5	2.0	1.0	3.0	1.5
3	1.5	1.0	0.0	1.5	0.0
4	1.5	0.0	-1.0	0.0	-1.5
5	1.5	-1.0	-2.0	-1.5	-5.0
6	1.5	-2.0	-3.0	-3.0	-4.5

The DATA file, including subject identification and item responses, was as follows. Note that in coding the item responses, a "1" indicated the "best" response and a "3" indicated the "poorest," as specified by the item difficulty parameters in IPOOL.

1	333211
2	332111
3	321212
4	112121
5	112233
6	222222
7	122221
8	322223
9	111333
10	111222
11	333222
12	333111
13	222111
14	211111

The following is an example of the logistic graded model (OPT4=1) with a 1.7 scaling factor. In this example the  $b$  parameters for the items were taken from columns 3-4 of the IPOOL file. The option and format cards for this example were

```

      6      0201 1 1.70
      (11,1X,F3.1,2(1X,F4.1))
222222
      1      2      3      4      5      6
000000
      (2A7,A1,6I1)
EXAMPLE RUN OF THE LOGISTIC GRADED MODEL--USES THE FIRST PAIR OF B
PARAMETERS FROM ITEM POOL

```

The output from this run was as follows:

LINPSCO

LINEAR POLYCHOTOMUS SCORING WITH TWO PARAMETER MODELS

PSYCHOMETRICS METHODS PROGRAM  
DEPARTMENT OF PSYCHOLOGY  
UNIVERSITY OF MINNESOTA  
MPLS. MINN. 55455

INUP = 6  
MMAX = 6  
IOMIT = 4  
OPT1 = 0  
OPT2 = 1  
OPT4 = 1  
MAXCAT = 2  
U = 1.7  
VARIABLE FORMAT FOR POOL = (I1,1X,F3.1,2(1X,F4.1))  
VARIABLE FORMAT FOR DATA = (2A7,A1,6I1)  
EXAMPLE RUN OF THE LOGISTIC GRADED MODEL--USES THE FIRST PAIR OF B  
PARAMETERS FROM ITEM POOL

ITEM ID = 1 REJECTION = 0  
A: 1.50  
B: 3.00 2.00

ITEM ID = 2 REJECTION = 0  
A: 1.50  
B: 2.00 1.00

ITEM ID = 3 REJECTION = 0  
A: 1.50  
B: 1.00 0

ITEM ID = 4 REJECTION = 0  
A: 1.50  
B: 0 -1.00

ITEM ID = 5 REJECTION = 0  
A: 1.50  
B: -1.00 -2.00

ITEM ID = 6 REJECTION = 0  
A: 1.50  
B: -2.00 -3.00

CASES READ = 14 CASES NOT CONVERGED = 0

Testee Identification	Proportion of "Best" Responses (Coded 1)	Maximum Likelihood Estimate of $\theta$	Response Pattern Information	Number of Items Used to Estimate $\theta$	Number of Iterations	Test Information Associated with $\hat{\theta}$	Estimated Standard Error of Measurement
1	.33	-.50	4.72	6	2	4.25	.46
2	.50	.50	4.72	6	2	4.25	.46
3	.33	.35	3.69	6	3	4.28	.52
4	.67	1.52	2.63	6	3	4.11	.62
5	.33	-.00	4.21	6	3	4.41	.49
6	.50	-.00	5.16	6	2	4.41	.44
7	.50	.51	4.88	6	2	4.25	.45
8	.50	-.51	4.88	6	2	4.25	.45
9	.50	-.00	.96	6	3	4.41	1.02
10	.50	.52	2.65	6	2	4.25	.61
11	.50	-1.51	4.86	6	2	4.11	.45
12	.50	-.00	4.20	6	3	4.41	.49
13	.50	1.51	4.86	6	2	4.11	.45
14	.83	2.69	3.12	6	2	2.67	.57

Following is an example of use of the normal ogive graded model (OPT4=2) using the same DATA and IPOOL as the previous example. In this example the  $b$  parameters for the items were taken from columns 5 and 6 of the IPOOL file. Input control cards for this example were

```

      6      0201 2      4
      (11,1X,F3.1,11X,F4.1,1X,F4.1)
      222222
      1      2      3      4      5      6
      000000
      (2A7,A1,6I1)
      EXAMPLE RUN OF THE NORMAL OGIVE GRADED MODEL--USES SECOND PAIR OF B
      PARAMETERS FROM ITEM POOL
  
```



Output was as follows:

LINPSCO

LINEAR POLYCHOTOMUS SCORING WITH TWO PARAMETER MODELS

PSYCHOMETRICS METHODS PROGRAM  
DEPARTMENT OF PSYCHOLOGY  
UNIVERSITY OF MINNESOTA  
MPLS. MINN. 55455

INUP     =     6  
MMAX     =     6  
IOMIT    =     4  
OPT1     =     0  
OPT2     =     1  
OPT4     =     2  
MAXCAT   =     2  
VARIABLE FORMAT FOR POOL =(I1,1X,F3.1,11X,F4.1,1X,F4.1)  
VARIABLE FORMAT FOR DATA=(2A7,A1,6I1)  
EXAMPLE RUN OF THE NORMAL OGIVE GRADED MODEL--USES SECOND PAIR OF B  
PARAMETERS FROM ITEM POOL

ITEM ID =     1     REJECTION = 0  
A:     1.50  
B:     4.50   3.00

ITEM ID =     2     REJECTION = 0  
A:     1.50  
B:     3.00   1.50

ITEM ID =     3     REJECTION = 0  
A:     1.50  
B:     1.50   0

ITEM ID =     4     REJECTION = 0  
A:     1.50  
B:     0   -1.50

ITEM ID =     5     REJECTION = 0  
A:     1.50  
B:    -1.50 -3.00

ITEM ID =     6     REJECTION = 0  
A:     1.50  
B:    -3.00 -4.50

CASES READ =   14   CASES NOT CONVERGED =   0

1	.33	-.75	2.97	6	2	3.36	.58
2	.50	.75	2.97	6	2	3.36	.58
3	.33	-.04	8.00	6	2	3.42	.35
4	.67	1.85	8.05	6	2	3.39	.35
5	.33	-.00	11.98	6	2	3.42	.29
6	0	-.00	11.58	6	2	3.42	.29
7	.33	1.09	9.50	6	2	3.39	.32
8	0	-1.09	9.50	6	2	3.39	.32
9	.50	-.00	12.80	6	1	3.42	.28
10	.50	.75	12.29	6	2	3.36	.29
11	0	-2.25	5.18	6	2	3.35	.44
12	.50	-.00	3.19	6	2	3.42	.56
13	.50	2.25	5.13	6	2	3.35	.44
14	.33	3.92	2.05	6	2	2.25	.70

Nominal logistic model. Following is an example of use of the nominal logistic model (OPT4=3). The DATA file was

```

10001 38 9946454134111122442211121114111141111121111111111
10002 38103834541114411414114411111411111411141114111
10003 38 94000542414112111441214121114111144111114111241
10005 381042450424222444424211142442421222421212121222241
10004 38 9954344414144441414444441411144411114114141144411
10005 381045285411144151214551121122111131111155555555551
10004 381055445443511111514111144111511114114111111114111
10008 38 99334942324551432331435241555555555555555555555
10009 3810534444142124152232134413124214124221114321222554
10010 381051509444444111444511141111411414111111111114141
10011 3810529454111111151141111111111111111111111111111
10012 3810584924423231334333114222251313251122144511242151
10013 3810540444414444144111441444111441411441114241511111
10015 38 9524554144512132544411411111111121241134411112141
10014 3810598354211241142144144441211214121141114111154411
10015 3810512954211141112441111411111111111111111111141
10014 381059528524413211314211142512111111411114111131451
10018 3810524844114411141324111414114244431441114114115141
10019 3810528254414411144241112433113442411241121311114114
10020 25104114344444444141441114444441414444141114144144444

```

IPOOL was

3417	0.000000	0.000000	0.000000	0.000000
3422	0.000000	0.000000	0.000000	0.000000
3425	0.870169	0.165559	-1.212113	0.176405
3431	0.842205	0.266821	-2.116617	1.006591
3434	1.032257	0.007821	-1.157598	0.110520
3437	0.900771	-0.711543	-1.432918	1.237689
3440	0.870227	0.313705	-1.132723	-0.059209
3443	0.510097	-0.183194	-1.493435	1.106532
3446	0.842335	-0.056809	-0.936221	0.143745
3449	0.947914	-0.510553	-1.591918	1.154556
3452	0.997044	-0.057076	-1.046348	0.106360
3455	0.220949	-0.511724	-0.842978	1.130754
3458	0.900633	0.452405	-1.315750	-0.040265
3461	1.324394	0.196605	-2.289410	0.768330
3464	0.935577	0.125247	-1.013213	-0.047610
3467	1.754989	-0.277927	-1.862908	0.405916
3470	1.267512	0.165917	-1.188908	-0.244462
3473	0.720900	-0.503622	-1.456565	1.237267
3476	0.400782	0.236527	-0.675647	0.032338
3479	-0.165035	-0.081050	-0.762658	0.968744
3482	0.914800	-0.087146	-0.807633	-0.020027
3485	0.880125	-0.561450	-1.133896	0.019221
3488	0.000000	0.000000	0.000000	0.000000
3491	0.000000	0.000000	0.000000	0.000000
3494	1.210009	-0.160504	-1.049275	-0.005201
3497	0.700199	-0.472306	-1.271001	1.040191
3500	0.941547	0.120504	-1.216024	0.154792
3503	1.010184	-0.227101	-2.405601	1.014638
3506	1.190701	0.289311	-1.692915	0.204903
3509	2.322618	-0.227105	-3.302494	1.207041
3512	1.382173	-0.138215	-1.203993	-0.039965
3515	1.750215	-0.061300	-2.359620	0.960275
3518	1.160829	-0.009702	-1.500254	0.349206
3521	0.157054	-0.506044	-1.636365	1.987376
3524	1.234110	-0.375024	-0.934679	-0.223813
3527	1.111208	0.312124	-1.527765	0.404369
3530	1.020419	0.034121	-0.950948	-0.106622
3533	0.550335	-0.287405	-1.247386	0.981450
3536	1.600121	-0.008501	-1.228285	-0.303305
3539	2.097420	-0.093601	-2.469919	0.460094
3542	1.591045	-0.042903	-1.013761	-0.334310
3545	1.080738	-0.088700	-1.734153	0.139205
3548	1.022427	0.153605	-1.046231	-0.159844
3551	1.574805	-0.443101	-1.656987	0.525223
3554	0.994791	-0.067707	-1.062704	0.135620
3557	1.010407	-0.057804	-1.145208	0.760626
3560	1.242329	0.078505	-1.435152	0.114271
3563	2.364325	-0.059700	-2.638202	1.133657
3566	1.121352	0.184404	-0.962735	-0.343020
3569	1.020360	-0.262202	-1.632686	0.274608
3572	1.350141	0.216193	-1.205470	-0.366864
3575	2.260497	-0.343707	-2.311491	0.389761
3578	0.000197	0.002103	-0.012300	0.007000
3581	0.970607	-0.472700	-1.024609	0.523197
3584	2.042249	0.393202	-1.696127	-0.757373
3587	0.700431	0.132907	-3.787033	-0.054433
3590	1.577801	0.151890	-1.134625	-0.373132
3593	2.120020	0.251712	-2.499535	0.094737
3596	1.540139	0.286402	-2.081177	0.249600
3599	2.042910	0.001304	-3.331305	1.207001
3602	1.564642	-0.007945	-1.296770	-0.257927
3605	2.290293	0.021340	-2.644136	0.326495
3608	1.280960	0.227021	-0.889192	-0.023795
3611	0.010197	-0.447594	-2.181694	-0.339909
3614	1.050257	-0.089429	-0.664305	-0.293623
3617	1.620671	-0.009501	-1.288305	-0.033315
3620	1.450187	0.264101	-1.620217	-0.102162
3623	1.770329	-0.348500	-2.527377	1.184546
3626	1.110323	-0.003904	-0.823243	-0.292120
3629	1.270163	-0.303637	-1.057345	0.081860
3632	1.330099	0.035376	-1.773168	0.400713
3635	2.390180	-0.114600	-3.584400	1.336280
3638	1.480476	-0.073010	-0.823643	-0.589715
3641	2.230098	-0.331490	-1.535212	-0.371390
3644	1.200460	-0.157705	-0.924427	-0.120298
3647	1.770003	-0.425000	-1.844767	0.313329
3650	1.000538	-0.131202	-0.860200	-0.069056
3653	1.630196	-0.343905	-1.242548	0.453307
3656	1.320054	0.287821	-1.959475	0.243501
3659	1.000725	0.560903	-3.220217	0.994529
3662	1.000690	-0.074201	-1.264006	-0.268341
3665	2.077034	-0.045201	-2.330619	0.930460
3668	0.691561	0.155307	-1.158516	0.311618
3671	1.310654	-1.378577	-1.585505	1.053456
3674	1.091246	0.129610	-1.359014	-0.460943
3677	3.629455	-0.593270	-3.131603	0.095448



```

INUP      = 43
MMAX      = 22
IOMI1     = 5
OPT1      = 0
OPT2      = 0
OPT4      = 3
MAXCAT    = 3
VARIABLE FORMAT FOR POOL =(I4,4(1X,F9.6),/,4X,4(1X,F9.6))
VARIABLE FORMAT FOR DATA=(2A7,A1,43I1)
EXAMPLE RUN OF THE NOMINAL LOGISTIC MODEL

```

ITEM ID = 0 0 REJECTION = 1  
A: 0 0 0 0  
B: 0 0 0 0

ITEM ID = 3251 REJECTION = 0  
A: 1.04 .01 -1.16 .11  
B: .91 -.71 -1.43 1.24

ITEM ID = 3422 REJECTION = 0  
A: .85 -.06 -.94 .14  
B: .95 -.51 -1.59 1.15

ITEM ID = 3421 REJECTION = 0  
A: .90 .45 -1.32 -.04  
B: 1.32 .20 -2.29 .77

ITEM ID = 3277 REJECTION = 0  
A: .94 .13 -1.01 -.05  
B: 1.75 -.28 -1.88 .41

ITEM ID = 3400 REJECTION = 0  
A: 1.27 .17 -1.19 -.24  
B: .72 -.50 -1.46 1.24

ITEM ID = 0 0 REJECTION = 1  
A: 0 0 0 0  
B: 0 0 0 0

ITEM ID = 3405 REJECTION = 0  
A: 1.22 -.16 -1.05 -.01  
B: .70 -.47 -1.27 1.04

ITEM ID = 3213 REJECTION = 0  
A: .94 .12 -1.22 .15  
B: 1.62 -.23 -2.41 1.01

ITEM ID = 3079 REJECTION = 0  
A: 1.38 -.14 -1.20 -.04  
B: 1.75 -.66 -2.06 .97

ITEM ID = 3002 REJECTION = 0  
A: 1.16 -.01 -1.50 .35  
B: .16 -.51 -1.64 1.99

ITEM ID = 3210 REJECTION = 0  
A: 1.38 -.04 -1.61 -.33  
B: 1.68 -.09 -1.73 .14

ITEM ID = 3404 REJECTION = 0  
A: .99 -.07 -1.06 .14  
B: 1.02 -.66 -1.15 .79

ITEM ID = 3400 REJECTION = 0  
A: 1.24 .08 -1.44 .11  
B: 2.36 -.86 -2.64 1.13

ITEM ID = 3021 REJECTION = 0  
A: 2.04 .39 -1.70 -.74  
B: 3.71 .13 -3.79 -.05

ITEM ID = 3221 REJECTION = 0  
A: 1.38 .13 -1.15 -.38  
B: 2.12 .28 -2.50 .09

ITEM ID = 3076 REJECTION = 0  
A: 1.29 .23 -.69 -.62  
B: 3.02 -.45 -2.18 -.39

ITEM ID = 3200 REJECTION = 0  
A: 1.20 -.16 -.92 -.12  
B: 1.77 -.24 -1.04 .31

ITEM ID = 3029 REJECTION = 0  
A: .69 .16 -1.16 .31  
B: 1.31 -1.38 -1.59 1.65

ITEM ID = 3078 REJECTION = 0  
A: 1.69 .13 -1.36 -.46  
B: 3.63 -.59 -3.13 .10

ITEM ID = 3209 REJECTION = 0  
A: 1.33 .04 -1.77 .41  
B: 2.39 -.11 -3.56 1.31

ITEM ID = 3023 REJECTION = 0  
A: 1.48 .07 -.82 -.59  
B: 2.24 -.33 -1.54 -.37

CASES READ = 20 CASES NOT CONVERGED = 0

Testee Identification		Proportion of "Best" Responses	Maximum Likelihood Estimate of $\theta$	Response Pattern Information	Number of Items Used to Estimate $\theta$	Number of Iterations	Test Information Associated with $\theta$	Estimated Standard Error of Measurement
8 99484	5	.50	-.63	9.16	20	3	9.12	.33
8103834	5	.65	-.02	6.24	20	3	6.21	.40
8 94000	5	.55	-.33	7.94	20	2	7.60	.35
8104245	0	.15	-1.31	12.19	20	2	12.11	.29
8 99543	4	.35	-1.10	11.43	20	2	11.44	.30
8104528	5	.59	-.47	7.11	17	2	7.11	.37
8105544	5	.71	-.07	5.17	17	3	5.16	.44
8 99334	9	.19	-1.73	9.55	16	2	9.55	.32
8105344	4	.32	-1.40	11.89	19	2	11.89	.29
8105150	9	.47	-.69	9.02	19	3	9.00	.33
8105294	5	.95	3.12	.59	19	3	.59	1.30
8103849	2	.16	-1.90	10.86	19	3	10.85	.30
8105404	4	.45	-.72	9.60	20	3	9.59	.32
8 95245	5	.56	-.51	7.98	18	3	7.97	.35
8105983	5	.40	-1.05	11.34	20	2	11.24	.30
8105129	5	.70	.00	6.17	20	3	6.14	.40
8105952	8	.47	-.85	10.09	19	3	10.09	.31
8105248	4	.55	-.49	8.39	20	3	8.39	.35
8105282	5	.40	-1.03	11.18	20	3	11.18	.30
5104114	3	.30	-.98	10.94	20	3	10.95	.30



APPENDIX C  
LINDSCO FORTRAN PROGRAM LISTING

```

PROGRAM LINDSCO (INPUT,OUTPUT,DATA,IPOOL,TAPE1=DATA,TAPE2=IPOOL,TA
1PE3,PUNCH)
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033

```

```

*
*IN THE NEXT DO LOOP THE ITEM PARAMETERS CORRESPONDING TO THE ITEMS
*IN INAD ARE RETRIEVED FROM A,B,C, AND LOADED INTO ADM,BDM,CDM RESP.
*THE ENTRIES IN THE ADM,BDM,CDM ARE ZEROED FOR THE CASE OF ZERO ITEM
*ID IN THE INAD
*
*****
DO 8 J=1,M
  IF (INAD(J).NE.0) GO TO 6
  ADM(J)=BDM(J)=CDM(J)=0
  GO TO 8
6  CONTINUE
  IFOUND=0
  DO 7 I=1,INUP
    IF (INAD(J).NE.ITEM(I)) GO TO 7
    IFOUND=1
    ADM(J)=A(I)
    BDM(J)=B(I)
    CDM(J)=C(I)
    GO TO 8
7  CONTINUE
  IF (IFOUND.EQ.0) INAD(J)=0
8  CONTINUE
  GO TO 11
C    THE NEXT SECTION IS USED IF OPTION 2 IS ON, IT WILL TAKE THE FI
C    PARAMETERS FROM THE POOL WITHOUT MAKING USE OF INAD
9  DO 10 I=1,M
    ADM(I)=A(I)
    BDM(I)=B(I)
    CDM(I)=C(I)
10  CONTINUE
C    IF OPTION 3 IS ON THE PARAMETERS A,B,C ARE CONSTRAINED WITHIN B
C    OF AMAX,AMIN,BMAX,BMIN,CMAX
11  IF (OPT3.EQ.0) GO TO 13
    DO 12 I=1,M
      IF (INAD(I).EQ.0) GO TO 12
      IF ((ADM(I).GT.AMAX).OR.(OPT3.EQ.3)) ADM(I)=AMAX
      IF (BDM(I).LT.BMIN) BDM(I)=BMIN
      IF (BDM(I).GT.BMAX) BDM(I)=BMAX
      IF ((CDM(I).GT.CMAX).OR.(OPT3.EQ.2)) CDM(I)=CMAX
12  CONTINUE
13  IF (OPT1.NE.1) GO TO 15
    DO 14 I=1,M
      PUNCH 57, INAD(I),KEY(I),IREJ(I),ADM(I),BDM(I),CDM(I)
14  CONTINUE
15  PRINT 57, (INAD(I),KEY(I),IREJ(I),ADM(I),BDM(I),CDM(I),I=1,M)
    PRINT 49
*****
*
*
*READ A SUBJECT FROM TAPE1 CALCULATE THETA, LOOP BACK TO 5 ETC.
*
*****
15  READ (1,IFORM) NAME,ID,(IRAW(I),I=1,M)
    DO 17 JJ=1,M
17  IRESP(JJ)=0
    IO=UNIT(1)
C    CHECK THE END OF FILE ON DATA FILE
    IF (IO.LE.3) GO TO 18
    PRINT 59, IO
18  IF (IO.EQ.0) GO TO 47
    N=N+1
    ITOT=0.0
C    SET THE ITEM ID TO ZERO FOR THE OMISSIONS (THAT IS READ IN FRO

```

C		133
C	AND SET THE RESPONSE VECTOR TO 1 IF THE ANSWER IS CORRECT	134
C		135
	DO 21 I=1,M	136
	INADS(I)=INAD(I)	137
	IF (IRAW(I).EQ.IOMIT) 19,20	138
19	INADS(I)=0	139
	GO TO 21	140
20	IF ((IRAW(I).EQ.KEY(I)).AND.(IREJ(I).EQ.0)) IRESP(I)=1	141
	ITOT=ITOT+FLOAT(IRESP(I))	142
21	CONTINUE	143
	ITINADS=0	144
	DO 22 KK=1,M	145
22	IF (INADS(KK).EQ.0) ITINADS=ITINADS+1	146
	KL=M-ITINADS	147
	ITOT=ITOT/FLOAT((KL))	148
	IF (ITOT.EQ.0.) 23,24	149
23	PRINT 60, NAME, ID	150
	IF (OPT4.EQ.3) GO TO 26	151
	T=-10.0	152
	SFORM=0.0	153
	TINFO=0.0	154
	IKL=KL	155
	ITER=0	156
	EXPTOT=0.0	157
	WRITE (3,63) NAME, ID, ITOT, T, SFORM, IKL, TINFO, EXPTOT, ITER	158
	GO TO 32	159
24	IF (ITOT.EQ.1.) 25,26	160
25	PRINT 61, NAME, ID	161
	IF (OPT4.EQ.3) GO TO 26	162
	T=10.0	163
	SFORM=0.0	164
	TINFO=0.0	165
	IKL=KL	166
	ITER=0	167
	EXPTOT=0.0	168
	WRITE (3,63) NAME, ID, ITOT, T, SFORM, IKL, TINFO, EXPTOT, ITER	169
	GO TO 32	170
	*****	171
	*	172
	*	173
	* NOW THE DATA IS READY TO MAKE THE CALLS TO THE APPROPRIATE ROUTINE	174
	* TO ESTIMATE THE THETA. OPTION 4 WILL DETERMINE THE METHOD BY WHICH	175
	* THE THETA ESTIMATE WILL BE FOUND	176
	*	177
	*	178
	*****	179
26	IKL=KL	180
	IF (OPT4-2) 27,28,29	181
27	CALL MAXLNO (IRESP, INADS, M, M, ADM, RDM, CDM, 50, .005, T, SFORM, IFAIL, TINFO, EXPTOT, ITER, SEM)	182
	GO TO 30	183
28	CALL MAXLK (M, INADS, IRESP, M, ADM, RDM, CDM, 50, .005, IFAIL, SFORM, T, TINFO, EXPTOT, ITER, SEM)	184
	GO TO 30	185
29	T=TS	186
	SFORM=TS	187
	SEM=0.0	188
	CALL HAYES (M, INADS, IRESP, M, ADM, RDM, CDM, T, SFORM, TINFO, EXPTOT, ITER)	189
	GO TO 31	190
30	CONTINUE	191
	IF (IFAIL.EQ.0) GO TO 31	192
	PRINT 62, NAME, ID	193
	SFORM AND T ARE SET TO -99.99 IN MAXLK IF NOT CONV	194
	NC=NC+1	195
	*****	196
		197
		198



```

*
*WRITE THE SUCCESSFUL RESULTS TO THE FILE TAPE3
*
*****
31  WRITE (3,63) NAME,ID,ITOT,T,SFORM,IKL,TINFO,EXPTOT,ITER,SEM
C      IF THERE IS NO SUBSCALE CALCULATIONS LOOP BACK TO READ A SUBJECT
C      OTHERWISE CONTINUE WITH THE COMPUTATIONS FOR SUBSCALES
32  IF (NSSC.EQ.0) GO TO 16
    DO 46 I=1,NSSC
      NI=NISS(I,1)
      ITOT=0.0
      DO 32 J=1,M
        ISADS(J)=INADS(J)
      DO 34 K=1,NI
        IF (ISAD(K,I)-INADS(J)) 34,33,34
33    ITOT=ITOT+FLOAT(IRESP(J))
      GO TO 35
34    CONTINUE
      ISADS(J)=0
35    CONTINUE
      ITISADS=0
      DO 36 KK=1,M
36    IF (ISADS(KK).NE.0) ITISADS=ITISADS+1
      KL=ITISADS
      ITOT=ITOT/FLOAT(KL)
      IF (ITOT.EQ.0.0) 37,38
37    PRINT 66, NAME,ID,NISS(I,2)
      IF (OPT4.EQ.3) GO TO 40
      T=-10.00
      TINFO=EXPTOT=0.0
      ITER=0
      SFORM=0.0
      WRITE (3,64) NAME,ID,NISS(I,2),ITOT,T,SFORM,IKL,TINFO,EXPTOT,ITER
      GO TO 46
38    IF (ITOT.EQ.1.0) 39,40
39    PRINT 67, NAME,ID,NISS(I,2)
      IF (OPT4.EQ.3) GO TO 40
      T=10.0
      TINFO=EXPTOT=0.0
      ITER=0
      SFORM=0.0
      WRITE (3,64) NAME,ID,NISS(I,2),ITOT,T,SFORM,IKL,TINFO,EXPTOT,ITER
      GO TO 46
40    IKL=KL
      IF (OPT4-2) 41,42,43
41    CALL MAXLNQ (IRESP,ISADS,M,M,ADM,BDM,CDM,50,.01,T,SFORM,IFAIL,TINF
10,EXPTOT,ITER,SEM)
      GO TO 44
42    CALL MAXLK (M,ISADS,IRESP,M,ADM,BDM,CDM,50,.01,IFAIL,SFORM,T,TINFO
1,EXPTOT,ITER,SEM)
      GO TO 44
43    T=TS
      SFORM=TSS
      SEM=0.0
      CALL HAYES (M,ISADS,IRESP,M,ADM,BDM,CDM,T,SFORM,TINFO,EXPTOT,ITER)
      GO TO 45
44    IF (IFAIL.EQ.0) GO TO 45
      PRINT 68, NAME,ID,NISS(I,2)
45    WRITE (3,64) NAME,ID,NISS(I,2),ITOT,T,SFORM,IKL,TINFO,EXPTOT,ITER,
1SEM
46    CONTINUE
      GO TO 16
47    PRINT 65, N,NC
      STOP
C
C

```

C		265
48	FORMAT (I5)	266
49	FORMAT (1H1)	267
50	FORMAT (2I4,X,4I1,6F5.2,2X,I2)	268
51	FORMAT (16I5)	269
52	FORMAT (80I1)	270
53	FORMAT (8A10)	271
54	FORMAT (T50,*LINDSCO*,/,T50*====*,////////,T20,*LINEAR DICHOTOMUS	272
	1 SCORING WITH THREE PARAMETER MODELS*,////////,T40*PSYCHOMETRIC METHOD	273
	2S PROGRAM*,/,T40*DEPARTMENT OF PSYCHOLOGY*,/,T40*UNIVERSITY OF MIN	274
	3NESOTA*,/,T40*MPLS. MINN. 55455*,/,///,T20*INUP*,T27*==I5,/,T20,*M	275
	4MAX*,T27*==I5,/,T20,*IOMIT*,T27*==I5,/,T20*OPT1*,T27*==I5,/,T20*O	276
	5PT2*,T27*==I5,/,T20*OPT3*,T27*==I5,/,T20*OPT4*,T27*==I5,/,T20*TS*	277
	6,T27*==F5.2,/,T20*TSS*,T27*==F5.2,/,T20*AMAX*,T27*==F5.2,/,T20,*RM	278
	7AX*,T27,*==F5.2,/,T20*BMIN*,T27,*==F5.2,/,T20*CMAX*,T27,*==F5.2,/,	279
	4T20,*VARIABLE FORMAT FOR POOL=*8A10,/,T20,*VARIABLE FORMAT FOR DAT	280
	9A=*8A10,/,T20,8A10,/,T20,8A10,/,T20,8A10,/,)	281
55	FORMAT (8A10)	282
56	FORMAT (2I5)	283
57	FORMAT (*1*,29X,*ITEM ID S*2X,*KEYS*2X,*REJECTIONS*,4X,*A*,9X,*B*,	284
	19X,*C*,/,60(/30X,I5,5X,I3,5X,I3,1X,F10.2,1X,F10.2))	285
58	FORMAT (///,40X,*ITEMS IN SUBSCALE NO=*I3,/,10(20I6,/) )	286
59	FORMAT (10X,*PARITY ERROR ON TAPE*,90X,I2)	287
60	FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS NO ANSWERS *,*CORRE	288
	1CT IN TOTAL SCALE*)	289
61	FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS ALL ANSWERS *,*CORRE	290
	1CT IN TOTAL SCALE*)	291
62	FORMAT (10X,*COMPUTATIONAL PROBLEMS WITH SUBJECT =*,2A10,* ID =*,A	292
	19,* IN TOTAL TE*)	293
63	FORMAT (X,2A10,A9,* T*,F5.2,2F7.2,I4,2F7.2,I4,F7.2)	294
64	FORMAT (X,2A10,A9,I2,F5.2,2F7.2,I4,2F7.2,I4,F7.2)	295
65	FORMAT (/10X,*CASES READ=*,I5,* CASES NOT CONVERGED=*,I5)	296
66	FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS NO ANSWERS *,*CORRE	297
	1CT IN SUBSCALE *,I5)	298
67	FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS ALL ANSWERS *,*CORRE	299
	1CT IN SUBSCALE *,I5)	300
68	FORMAT (10X,*MAXIMUM LIKELIHOOD ESTIMATION DOES NOT CONVERGE*,*FOR	301
	1 THE SUBJECT = *,2A10,* ID = *,A9,* ON SUBSCALE *,I5)	302
	END	303
	SUBROUTINE BAYES (M,ITM,RESP,N,A,B,C,BTHET,BVAR,TINFO,EXPTOT,ITER)	1
	INTEGER PESP(M),ITM(M)	2
	REAL A(N),B(N),C(N)	3
	DO 1 I=1,M	4
	IF (ITM(I).EQ.0) GO TO 1	5
	CALL BSCOR (BTHET,BVAR,B(I),A(I),C(I),RESP(I))	6
1	CONTINUE	7
	CALL NOSTAT (M,ITM,A,B,C,BTHET,TINFO,EXPTOT)	8
	ITER=0	9
	RETURN	10
	END	11
	SUBROUTINE BSCOR (BTHET,BVAR,DIF,DIS,GUESP,IRFSP)	1
	D=(DIF-BTHET)/SQRT(2.0*(1.0/DIS**2+BVAR))	2
	ERFD=ERFNP(D)	3
	EDSQ=EXP(D**2)	4
	IF (EDSQ.EQ.0.0) RETURN	5
	EDSQI=1.0/EDSQ	6
	XKINV=0.5*(1.0-ERFD)	7
	XLINV=GUESP*(1.0-GUESP)*XKINV	8
	IF ((XLINV.FQ.0.0).OR.(XKINV.EQ.0.0)) RETURN	9
	XL=1.0/XLINV	10
	IF (IKESP.NE.1) GO TO 1	11
	S=0.398942*(SQRT(BVAR)/SQRT(1.0+(1.0/DIS**2)/BVAR))*(1.0/XKINV)*ED	12
	1SQI	13
	T=1.0-1.772454*D*EDSQ*(1.0-ERFD)	14
	BTHET=BTHET+(1.0-GUESP)*XKINV*XL*S	15
	BVAR=BVAR-(1.0-GUESP)*XKINV*XL*S**2*(T-GUESP*XL)	16

```

1      RETURN
      HTHET=HTHET-0.797395*(BVAR/SQRT(1.0/DIS**2+BVAR))*EDSQI*(1.0/(1.0+
17      1ERFD))
18      PART1=1.128379/(1.0+(1.0/DIS**2)*(1.0/BVAR))
19      PART2=1.0/(EDSQ*(1.0+ERFD))**2
20      PART3=0.564190+0*EDSQ*(1.0+ERFD)
21      BVAR=BVAR*(1.0-PART1*PART2*PART3)
22      RETURN
23      END
24      REAL FUNCTION ERFNP (X)
25      DATA A1/0.254330/
1      DATA A2/-0.284497/
2      DATA A3/1.421414/
3      DATA A4/-1.453152/
4      DATA A5/1.061405/
5      DATA P/0.327591/
6      ERFNP=0.0
7      IF (X.EQ.0.0) RETURN
8      S=SIGN(1.0,X)
9      Y=ABS(X)
10     IF (Y.LT.6.0) GO TO 1
11     ERFNP=ES
12     RETURN
13     Y2=Y*Y
14     T=1.0/(1.0+P*Y)
15     AT=((A1+(A2+(A3+(A4+A5*T)*T)*T)*T)*T)
16     EAT=AT/EXP(Y2)
17     ERFNP=(1.0-EAT)*ES
18     RETURN
19     END
20     SUBROUTINE MAXLK (M,ITM,RESP,N,A,B,C,MAX,EPS,IFAIL,SDRV,THETA,TINF
21     10,EXPTOT,NUMITS,SEM)
22     EXTERNAL FDDLOG,SDDLOG
23     INTEGER RESP(M)
24     DIMENSION A(N), B(N), C(N), ITM(M)
25     C*** USES MAXIMUM LIKELIHOOD LOGISTIC SCORING ALGORITHM AND RESPONSE
26     C*** MODEL
27     C*** HISSECTION IS USED TO PROVIDE THE INITIAL GUESS FOR THE
28     C*** NEWTON-RAPHSON METHOD
29     CALL BISECT (FDDLOG,RESP,A,B,C,M,ITM,5,GUESS)
30     CALL NEWTRAP (FDDLOG,SDDLOG,RESP,A,B,C,M,ITM,MAX,EPS,NUMITS,GUESS,
31     1THETA,SDRV,IFAIL)
32     IF (IFAIL.EQ.1) 1,2
33     C*** NEWTON RAPHSON DID NOT CONVERGE
34     1 CALL NWTRR (THETA,SDRV,SEM,TINFO,EXPTOT)
35     RETURN
36     C
37     2 CALL LGSTAT (M,ITM,A,B,C,THETA,TINFO,EXPTOT)
38     SEM=1.0/SQRT(ABS(SDRV))
39     RETURN
40     END
41     FUNCTION FDDLOG (RESP,ITM,A,B,C,M,THETA)
42     INTEGER RESP(M),RIGHT
43     DIMENSION A(M), B(M), C(M), ITM(M)
44     DATA XMAX,XMIN/200.0,-200.0/
45     DATA D,RIGHT/1.7,1/
46     C*** CALCULATES FIRST DERIVATIVE OF LOG-LIKELIHOOD FUNCTION OF A
47     C*** RESPONSE VECTOR FOR THE LOGISTIC MODEL
48     SUM=C.0
49     DO 1 I=1,M
50     IF (ITM(I).EQ.0) GO TO 1
51     X=D*A(I)*(THETA-H(I))
52     IF (X.LT.XMIN) X=XMIN
53     IF (X.GT.XMAX) X=XMAX

```



```

EXF=EXP(X)
AE=A(I)*EXF
SUM=SUM-AE/(EXF+1.0)
IF (RESP(I).NE.RIGHT) GO TO 1
CE=C(I)+EXF
SUM=SUM+AE/CE
1 CONTINUE
FOOLOG=-1.7*SUM
RETURN
END
FUNCTION SDDLOG (RESP,ITM,A,B,C,M,THETA)
INTEGER RESP(M),RIGHT
DIMENSION ITM(M), A(M), B(M), C(M)
DATA XMAX,XMIN/200.0,-200.0/
DATA D,RIGHT/1.7,1/
C*** CALCULATES SECOND DERIVATIVE OF LOG-LIKELIHOOD FUNCTION
C*** OF A RESPONSE VECTOR FOR THE LOGISTIC MODEL
SUM=0.0
DO 1 I=1,M
IF (ITM(I).EQ.0) GO TO 1
X=0*A(I)*(THETA-B(I))
IF (X.LT.XMIN) X=XMIN
IF (X.GT.XMAX) X=XMAX
EXF=EXP(X)
AE=A(I)*EXF
SUM=SUM-A(I)*AE/((1.0+EXF)*(1.0+EXF))
IF (RESP(I).NE.RIGHT) GO TO 1
CE=C(I)+EXF
SUM=SUM+A(I)*C(I)*AE/(CE*CE)
1 CONTINUE
SDDLOG=-2.89*SUM
RETURN
END
SUBROUTINE BISECT (F1,RESP,A,B,C,M,ITM,NITER,BMID)
INTEGER RESP(M)
DIMENSION A(M), B(M), C(M), ITM(M)
C*** CALCULATES APPROXIMATE ROOT OF F1 BY BISECTION;
C*** BISECTING NITER (NUMBER OF ITERATIONS) TIMES.
C*** BMID IS BEST CURRENT GUESS AT ROOT THETA
C
C*** INITIALIZE LEFT BOUND AND F1(BOUND) AND RIGHT BOUND F1(BOUND)
BL=-5.0
BR=5.0
BMID=0.0
FL=F1(RESP,ITM,A,B,C,M,BL)
FR=F1(RESP,ITM,A,B,C,M,BR)
C*** TEST FOR NO ROOT IN INTERVAL--RETURN IF NO SOLUTION
IF ((TL*TR).GT.0.0) RETURN
C
C*** NOW CALCULATE BISECTIONS NITER TIMES
DO 3 I=1,NITER
TMID=F1(RESP,ITM,A,B,C,M,BMID)
IF ((TMID*TL).GT.0.0) GO TO 1
C*** REPLACE RIGHT BOUND WITH BMID
BR=BMID
GO TO 2
C*** REPLACE LEFT BOUND WITH BMID
1 TL=TMID
BL=BMID
C*** FIND NEW MIDPOINT BMID
2 BMID=(BL+BR)/2.0
3 CONTINUE
RETURN
END
SUBROUTINE NEWTRAP (F1,F2,RESP,A,B,C,M,ITM,NITER,EPS,NUMITS,GUESS,
1 THETA,SORV,IFAIL)
2

```

```

INTEGER RESP(M)
DIMENSION A(M), B(M), C(M), ITM(M)
C*** CALCULATES THE ROOT OF F1 GIVEN ITS FIRST DERIVATIVE F2
C*** AND AN INITIAL GUESS USING NEWTON-RAPHSON METHOD
C*** THETA IS APPR. TO THE ROOT: SDRV IS F2(THETA)
NUMITS=0
THETA=GUESS
C*** LOOP UNTIL ERR<EPS OR NUMBER OF ITERATIONS BECOMES TOO LARGE
1 FDRV=F1(RRESP,ITM,A,B,C,M,THETA)
SDRV=F2(RRESP,ITM,A,B,C,M,THETA)
ERR=FDRV/SDRV
THETA=THETA-ERR
NUMITS=NUMITS+1
C*** EXIT LOOP CRITERION
IF ((NUMITS.LT.NITER).AND.(ABS(ERR).GT.EPS)) GO TO 1
C*** END LOOP. TEST FOR CONVERGENCE AND SET IFAIL
IFAIL=0
IF (ABS(ERR).LT.EPS) RETURN
C
C*** NEWTON RAPHSON METHOD DOES NOT CONVERGE
IFAIL=1
RETURN
END
SUBROUTINE NWTRK (THETA,SFORM,SEM,TINFO,EXPTOT)
C*** SETS ERROR VALUES FOR THE CASE IN WHICH NEWTON RAPHSON FAILS
C*** TO CONVERGE
THETA=-99.99
SFORM=-99.99
SEM=-99.99
TINFO=-99.99
EXPTOT=-99.99
RETURN
END
SUBROUTINE LGSTAT (M,ITM,A,B,C,THETA,TINFO,EXPTOT)
DIMENSION A(M), B(M), C(M), ITM(M)
DATA XMAX,XMIN/12.0,-12.0/
TINFO=0.0
EXPTOT=0.0
KOUNT=0
C
DO 1 I=1,M
IF (ITM(I).EQ.0) GO TO 1
KOUNT=KOUNT+1
ARGU=-1.7*A(I)*(THETA-B(I))
IF (ARGU.GT.XMAX) ARGU=XMAX
IF (ARGU.LT.XMIN) ARGU=XMIN
P=C(I)+((1.0-C(I))*(1.0/(1.0+EXP(ARGU))))
Q=1.0-P
EARG=EXP(-ARGU)
PPRIME=EARG/((1.0+EARG)*(1.0+EARG))
PPRIME=PPRIME*(1.0-C(I))*A(I)*1.7
TINFO=TINFO+(PPRIME*PPRIME)/(P*Q)
EXPTOT=EXPTOT+P
1 CONTINUE
EXPTOT=EXPTOT/FLOAT(KOUNT)
RETURN
END
SUBROUTINE MAYLNO (RESP,ITM,M,N,A,B,C,MAX,EPS,THETA,SDRV,IFAIL,TIN
1FO,EXPTOT,NUMITS,SEM)
EXTERNAL FDRGV,SDRNV
INTEGER RESP(M)
DIMENSION ITM(N), A(N), B(N), C(N)
C*** USES MAXIMUM LIKELIHOOD NORMAL OGIVE SCORING ALGORITHM AND
C*** RESPONSE VECTOR
C*** BISECTION IS USED TO PROVIDE THE INITIAL GUESS FOR THE
C*** NEWTON RAPHSON METHOD

```

```

C      CALL BISECT (FONOGV,RESP,A,B,C,M,ITM,5,GUESS)
C      CALL NEWTRAP (FONOGV,SDNOGV,RESP,A,B,C,M,ITM,MAX,EPS,NUMITS,GUESS,
1THETA,SDRV,IFAIL)
      IF (IFAIL.EQ.1) 1,2
C***  NEWTON RAPHSON DID NOT CONVERGE
1      CALL MWERR (THETA,SDRV,SEM,TINFO,EXPTOT)
      RETURN
C
2      CALL NOSTAT (M,ITM,A,B,C,THETA,TINFO,EXPTOT)
      SDRV=ABS(SDRV)
      SEM=1.0/SQRT(SDRV)
      RETURN
END
FUNCTION FONOGV (RESP,ITM,A,B,C,M,THETA)
  INTEGER RESP(M),RIGHT
  DIMENSION A(M), B(M), C(M), ITM(M)
  DATA PI,RIGHT/3.141592,1/
  DATA XMAX,XMIN/7.0,-7.0/
C***  CALCULATES FIRST DERIVATIVE OF LOG-LIKELIHOOD FUNCTION OF
C***  A RESPONSE VECTOR FOR THE NORMAL OGIVE MODEL
C
      SUM=0.0
      ROOTPI=1.0/SQRT(2.0*PI)
      DO 2 I=1,M
      IF (ITM(I).EQ.0) GO TO 2
      TEMP=A(I)*(THETA-B(I))
      IF (TEMP.GT.XMAX) TEMP=XMAX
      IF (TEMP.LT.XMIN) TEMP=XMIN
      X=-(TEMP*TEMP)/2.0
      DENOM1=ROOTPI*A(I)*(1.0-C(I))*EXP(X)
      DENOM=C(I)+(1.0-C(I))*CDFN(TEMP)
      IF (RESP(I).EQ.RIGHT) GO TO 1
      DENOM=- (1.0-DENOM)
1      SUM=SUM+(DENOM1/DENOM)
2      CONTINUE
      FONOGV=SUM
      RETURN
END
FUNCTION SDNOGV (RESP,ITM,A,B,C,M,THETA)
  INTEGER RESP(M),RIGHT
  DIMENSION A(M), B(M), C(M), ITM(M)
  DATA PI,RIGHT/3.141592,1/
  DATA XMAX,XMIN/7.0,-7.0/
C***  CALCULATES SECOND DERIVATIVE OF LOG-LIKELIHOOD FUNCTION
C***  OF A RESPONSE VECTOR FOR THE NORMAL OGIVE MODEL
C
      SUM=0.0
      ROOTPI=1.0/SQRT(2.0*PI)
      DO 2 I=1,M
      IF (ITM(I).EQ.0) GO TO 2
      TEMP1=A(I)*(THETA-B(I))
      IF (TEMP1.GT.XMAX) TEMP1=XMAX
      IF (TEMP1.LT.XMIN) TEMP1=XMIN
      X=-TEMP1*TEMP1/2.0
      TEMP2=ROOTPI*(1.0-C(I))*A(I)*EXP(X)
      FIRNUM=TEMP2*TEMP2
      SECNUM=TEMP2*A(I)*TEMP1
      SDENOM=C(I)+(1.0-C(I))*CDFN(TEMP1)
      FSDENOM=SDENOM*SDENOM
      IF (RESP(I).EQ.RIGHT) GO TO 1
      FSDENOM=(1.0-SDENOM)*(1.0-SDENOM)
      SDENOM=- (1.0-SDENOM)
1      SUM=SUM- (FIRNUM/FSDENOM) - (SECNUM/SDENOM)
2      CONTINUE

```



	SNDGV=SUM	27
	RETURN	28
	END	29
	SUBROUTINE NCSTAT (M,ITM,A,H,C,THETA,TINFO,EXPTOT)	1
	DIMENSION A(M), B(M), C(M), ITM(M)	2
	DATA PI/3.1415927	3
	DATA XMAX,XMIN/7.0,-7.0/	4
	TINFO=0.0	5
	EXPTOT=0.0	6
	KOUNT=0	7
		8
	DO 1 I=1,M	9
	IF (ITM(I).EQ.0) GO TO 1	10
	KOUNT=KOUNT+1	11
	TEMP=A(I)*(TH-THETA-2(I))	12
	IF (TEMP.GT.XMAX) TEMP=XMAX	13
	IF (TEMP.LT.XMIN) TEMP=XMIN	14
	P=C(I)+(1.0-C(I))*CDFN(TEMP)	15
	Q=1.0-P	16
	TEMP=-TEMP*TEMP/2.0	17
	PPRIME=(1.0/SQRT(2.0*PI))*(1.0-C(I))*A(I)*EXP(TEMP)	18
	TINFO=TINFO+(PPRIME*PPRIME)/(P*Q)	19
	EXPTOT=EXPTOT+P	20
1	CONTINUE	21
	EXPTOT=EXPTOT/KOUNT	22
	RETURN	23
	END	24

APPENDIX D  
ADADSCO FORTRAN PROGRAM LISTING

```

PROGRAM ADADSCO (INPUT,OUTPUT,DATA,IPOOL,TAPE1=DATA,TAPE2=IPOOL,TA
1PE3)
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66

```

DIMENSION ITEM(600), A(600), B(600), C(600), KEY(600), IREJ(80), I  
 1FORM2(8), IRAN(80), INADS(80), IRESP(80), ADM(80), BDM(80), CDM(80  
 2), DESC(24), NAME(2), IFORM1(8)  
 INTEGER OPT1,OPT2,OPT3,OPT4  
 REAL ITOT  
 N=NC=0  
 C\*\*\*\*\*  
 C  
 C READ OPTIONS AND PROGRAM PARAMETER FROM INPUT FILE, DATA IS ON TAPE 2  
 C  
 C\*\*\*\*\*  
 IPOOL=2  
 READ 20, INUP,MMAX,OPT1,OPT2,OPT3,OPT4,TS,TSS,AMAX,BMIN,BMAX,CMAX,  
 1IFLAG,IOMIT  
 READ 22, (IFORM1(I),I=1,8)  
 C IFORM1 IS THE VARIABLE FORMAT FOR THE ITEM POOL  
 READ (IPOOL,IFORM1) (ITEM(I),A(I),B(I),C(I),KEY(I),I=1,INUP)  
 C\*\*\*\*\*  
 C  
 C  
 C START READING THE SPECIFIC DATA (SPECIFIC FOR THE RUN) FROM THE INPUT  
 C INAD IS THE ITEM IDS ADMINISTERED  
 C IREJ IS THE REJECTED ITEM IDS  
 C IRESP IS THE RESPONSE VECTOR  
 C  
 C  
 C\*\*\*\*\*  
 READ 21, MNUM,(IREJ(I),I=1,MNUM)  
 READ 22, (IFORM2(I),I=1,8)  
 C IFORM2 IS THE VARIABLE FORMAT FOR THE SUBJECT DATA  
 C  
 READ 24, DESC  
 PRINT 23, INUP,MMAX,IOMIT,IFLAG,OPT1,OPT2,OPT3,OPT4,TS,TSS,AMAX,BM  
 1AX,BMIN,CMAX,IFORM1,IFORM2,DESC  
 C\*\*\*\*\*  
 C  
 C  
 C READ A SUBJECT FROM TAPE 1 CALCULATE THETA, LOOP BACK TO 5 ETC.  
 C  
 C  
 C\*\*\*\*\*  
 1 READ (1,IFORM2) ID,NAME,M,(IRESP(I),I=1,MMAX),(INADS(I),I=1,MMAX)  
 M=MIN0(M,MMAX)  
 IO=UNIT(1)  
 C CHECK THE END OF FILE ON DATA FILE  
 C  
 IF (IO.LE.0) GO TO 2  
 PRINT 25, IO  
 2 IF (IO.EQ.0) GO TO 19  
 N=N+1  
 ITOT=0.0  
 C SET THE ITEM ID TO ZERO FOR THE OMISSIONS (THAT IS READ IN FRO  
 C  
 C AND SET THE RESPONSE VECTOR TO 1 IF THE ANSWER IS CORRECT  
 DO 5 I=1,M  
 C SET THE ITEM IDS IN IREJ TO ZERO  
 IF (MNUM.EQ.0) GO TO 4  
 DO 3 IJ=1,MNUM  
 IF (INADS(I).NE.IREJ(IJ)) GO TO 3  
 INADS(I)=0  
 GO TO 5  
 3 CONTINUE  
 4 CONTINUE  
 IF (IRESP(I).EQ.IOMIT) INADS(I)=0

```

      IF (INADS(I).EQ.0) GO TO 5
      CALL SEARCH (INUP,INADS(I),ADM(I),BDM(I),COM(I),KKE,ITEM,A,B,C,KEY
1, ID)
C      IF THE FLAG IFLAG IS NOT ZERO IT IS TAKEN TO BE THE DUMMY KEY
C      IF IT IS ZERO THEN KEY IS READ FROM THE POOL AN LEFT IN KKE
      IF (IFLAG.NE.0) KKE=IFLAG
      IRES=0
      IF (IKRESP(I).NE.0,KKE) IRFS=1
      IRESP(I)=IRFS
5      CONTINUE
C*****
C      *
C      *
CIN THE NEXT DO LOOP THE ITEM PARAMETERS CORRESPONDING TO THE ITEMS
CIN INAD ARE RETRIVED FROM A,B,C, AND LOADED INTO ADM,BDM,COM RESP.
C THE ENTRIES IN THE ADM,BDM,COM ARE ZEROED FOR THE CASE OF ZERO ITEM
CID IN THE INAD
C      *
C      *
C*****
C      IF OPTION 3 IS ON THE PARAMETERS A,B,C ARE CONSTRAINED WITHIN B
C      OF AMAX,AMIN,BMAX,BMIN,CMAX
      IF (OPT3.EQ.0) GO TO 7
      DO 6 I=1,M
      IF (INADS(I).EQ.0) GO TO 6
      IF ((ADM(I).GT.AMAX).OR.(OPT3.EQ.3)) ADM(I)=AMAX
      IF (BDM(I).LT.BMIN) BDM(I)=BMIN
      IF (BDM(I).GT.BMAX) BDM(I)=BMAX
      IF ((COM(I).GT.CMAX).OR.(OPT3.EQ.2)) COM(I)=CMAX
6      CONTINUE
C      OPTION 1 WILL PRINT THE SPECIFIC DATA IF ITS ON
C
7      CONTINUE
      IF ((OPT1.EQ.0).OR.(OPT1.GT.10)) GO TO 8
      OPT1=OPT1+1
      PRINT 26, ID, NAME, (INADS(I), IRESP(I), ADM(I), BDM(I), COM(I), II=
11, M)
8      CONTINUE
      ITINADS=0
      DO 9 KK=1,M
      ITOT=ITOT+IRESP(KK)
9      IF (INADS(KK).EQ.0) ITINADS=ITINADS+1
      KL=M-ITINADS
      ITOT=ITOT/FLOAT((KL))
      IF (ITOT.EQ.0) 10,11
10     PRINT 27, NAME, ID
      IF (OPT4.EQ.3) GO TO 13
      F=-10.0
      SFORM=0.0
      FINFO=0.0
      EXPTOT=0.0
      SEM=0.0
      ITER=0
      IKL=KL
      GO TO 18
11     IF (ITOT.EQ.1.0) 12,13
12     PRINT 31, NAME, ID
      IF (OPT4.EQ.3) GO TO 13
      F=10.0
      SFORM=0.0
      FINFO=0.0
      EXPTOT=0.0
      SEM=0.0
      ITER=0
      IKL=KL
      GO TO 18

```



```

C***** 133
C 134
C 135
C NOW THE DATA IS READY TO MAKE THE CALLS TO THE APPROPRIATE ROUTINE 136
C TO ESTIMATE THE THETA. OPTIN 4 WILL DETERMINE THE METHOD BY WHICH 137
C THE THETA ESTIMATE WILL BE FOUND 138
C 139
C 140
C***** 141
13 IKL=KL 142
   IF (OPT4-2) 14,15,16 143
14 CALL MAXLNO (IRES,INADS,M,M,ADM,BDM,COM,50,.005,T,SFORM,IFAIL,TIN 144
   INFO,EXPTOT,ITER,SEM) 145
   GO TO 17 146
15 CALL MAXLK (M,INADS,IRES,M,ADM,BDM,COM,50,.005,IFAIL,SFORM,T,TINF 147
   INFO,EXPTOT,ITER,SEM) 148
C 149
   GO TO 17 150
16 T=TS 151
   SFORM=TSS 152
   CALL BAYES (M,INADS,IRES,M,ADM,BDM,COM,T,SFORM,TINFO,EXPTOT) 153
   ITER=0 154
   SEM=0.0 155
   GO TO 18 156
17 IF (IFAIL.EQ.0) GO TO 18 157
   PRINT 28, NAME,ID 158
C SFORM AND T ARE SET TO -99.99 IN MAXLK IF NOT CONV 159
   NC=NC+1 160
C***** 161
C 162
CWRITE THE SUCCESSFUL RESULTS TO THE FILE TAPE3 163
C 164
C***** 165
18 WRITE (3,29) NAME,ID,ITOT,T,SFORM,IKL,TINFO,EXPTOT,ITER,SEM 166
   GO TO 1 167
19 PRINT 30, N,NC 168
C 169
   STOP 170
C 171
C 172
20 FORMAT (2I4,X,4I1,6F5.2,I2,I2) 173
21 FORMAT (16I5) 174
22 FORMAT (8A10) 175
23 FORMAT (T50,*ADADSCO*,/,T50*====*,////////,T20,*ADAPTIVE DICHOTOM 176
   IUS SCORING WITH THREE PARAMETER MODELS*,////,T40*PSYCHOMETRIC METH 177
   ODS PROGRAM*,/,T40*DEPARTMENT OF PSYCHOLOGY*,/,T40*UNIVERSITY OF M 178
   INNESOTA*,/,T40*MPLS. MINN. 55455*,/,T20*INUP*,T27*=*I5,/,T20, 179
   4*MMAX*,T27*=*I5,/,T20,*IDMIT*,T27*=*I5,/,T20*IFLAG*,T27*=*I5,/,T 180
   520*OPT1*,T27*=*I5,/,T20*OPT2*,T27*=*I5,/,T20*OPT3*,T27*=*I5,/,T20* 181
   60PT4*,T27*=*I5,/,T20*IS*,T27*=*F5.2,/,T20*TSS*,T27*=*F5.2,/,T20*A 182
   7MAX*,T27*=*F5.2,/,T20,*RMAX*,T27,*=*F5.2,/,T20*PMIN*,T27,*=*F5.2, 183
   *,T20*CMAX*,T27,*=*F5.2,/,T20,*VARIABLE FORMAT FOR POOL=*8A10,/,T20 184
   9,*VARIABLE FORMAT FOR DATA=*8A10,/,T20,8A10,/,T20,8A10,/,T20,8A10, 185
   F/) 186
24 FORMAT (8A10) 187
25 FORMAT (10X,*PARITY ERROR ON TAPE*,90X,I2) 188
26 FORMAT (10X,A9,2A10,/, (1X,I4,2X,I2,2X,3F10.2)) 189
27 FORMAT (10X,* SUBJECT *,2A10,*ID= *,A9,*HAS NO RIGHT ANSWERS*) 190
28 FORMAT (10X,* MAXIMUM LIKELIHOOD ESTIMATION DOES NOT CONVERGE*,* F 191
   OR THE SUBJECT = *,2A10,* ID= *,A9) 192
29 FORMAT (X,2A10,A9,F5.2,2F7.2,I4,2F7.2,I4,F7.2) 193
30 FORMAT (10X,* CASES READ=*,I5,* CASES NOT CONVERGED=*,I5) 194
31 FORMAT (10X,* SUBJECT *,2A10,*ID= *,A9,*HAS ALL ANSWERS RIGHT*) 195
   END 196
SUBROUTINE SEARCH (INUP,ID,A,B,C,KEY,ITM,AP,BP,CP,KEYP,IDNU4) 1
DIMENSION AP(INUP), BP(INUP), CP(INUP), KEYP(INUP), ITM(INUP) 2

```

```

      INTEGER FLAG
      DO 1 I=1,INUP
      IF (ID.NE.ITM(I)) GO TO 1
      A=AP(I)
      B=BP(I)
      C=CP(I)
      KEY=KEYP(I)
      CALL PCHECK (A,B,C,ID,INUM)
      RETURN
1     CONTINUE
      PRINT 2, ID,INUM
      ID=J
      RETURN
C
C
C
2     FORMAT (10X,* ITEM =*,I4,* IS NOT IN THE POOL FOR SUBJECT ID =*,A9
1)
      END
      SUBROUTINE PCHECK (A,B,C,ID,INUM)
C*** CHECK WHETHER OR NOT ITEM PARAMETERS ARE VALID
C*** IF NOT, ERROR MESSAGE IS PRINTED
C
      IF (A.LE.0.0) PRINT 1, ID,A,INUM
      IF ((-5.0.GT.B).OR.(B.GT.5.0)) PRINT 2, ID,B,INUM
      IF ((0.0.GT.C).OR.(C.GT.1.0)) PRINT 3, ID,C,INUM
      RETURN
C
C
C
1     FORMAT (10X,*ITEM =*,I4,* HAS THE INVALID A PARAMETER OF *,F5.2,*
1     A MUST BE GREATER THAN 0.0*,/10X,*ERROR FOUND *,*FOR THE SUBJECT
2 WITH ID =*,A9)
2     FORMAT (10X,*ITEM =*,I4,* HAS THE EXTREME B PARAMETER OF *,F5.2,5X
1,*ERROR FOUND FOR THE SUBJECT WITH ID =*,A9)
3     FORMAT (10X,*ITEM =*,I4,* HAS THE INVALID C PARAMETER OF *,F5.2,*
1     C MUST BE BETWEEN 0.0 AND 1.0*,/10X,*ERROR FOUND *,*FOR THE SUBJ
2ECT WITH ID =*,A9)
      END
      SUBROUTINE BAYES (M,ITM,RESP,N,A,B,C,BTHET,BVAR,TINFO,EXPTOT)
      INTEGER RESP(M),ITM(M)
      REAL A(N),B(N),C(N)
      DO 1 I=1,M
      IF (ITM(I).EQ.0) GO TO 1
      CALL BSCOR (BTHET,BVAR,I(I),A(I),C(I),RESP(I))
1     CONTINUE
      CALL NOSTAT (M,ITM,A,B,C,BTHET,TINFO,EXPTOT)
      RETURN
      END
      SUBROUTINE BSCOR (BTHET,BVAR,DIF,DIS,GUESP,IRESP)
      D=(DIF-BTHET)/SQRT(2.0*(1.0/DIS**2+BVAR))
      ERFD=ERFNP(D)
      EDSQ=EXP(D**2)
      IF (EDSQ.EQ.0.0) RETURN
      EDSQ=1.0/EDSQ
      XKINV=0.5*(1.0-ERFD)
      XLINV=GUESP+(1.0-GUESP)*XKINV
      IF ((XLINV.EQ.0.0).OR.(XKINV.EQ.0.0)) RETURN
      XL=1.0/XLINV
      IF (IRESP.NE.1) GO TO 1
      S=0.398942*(SQRT(BVAR)/SQRT(1.0+(1.0/DIS**2)/BVAR))*(1.0/XKINV)*ED
1SQI
      T=1.0-1.72454*D*EDSQ*(1.0-ERFD)
      BTHET=BTHET+(1.0-GUESP)*XKINV*XL*S
      BVAR=BVAR-(1.0-GUESP)*XKINV*XL*S**2*(T-GUESP*XL)

```

```

1      RETURN
      BTHET=BTHET-0.797885*(BVAR/SQRT(1.0/DIS**2+BVAR))*EDSQI*(1.0/(1.0+
1ERFD))
      PART1=1.128379/(1.0+(1.0/DIS**2)*(1.0/BVAR))
      PART2=1.0/(EDSQ*(1.0+ERFD)**2
      PART3=0.564190+0*EDSQ*(1.0+ERFD)
      BVAR=BVAR*(1.0-PART1*PART2*PART3)
      RETURN
      END
      REAL FUNCTION ERFNP (X)
      DATA A1/0.254830/
      DATA A2/-0.284497/
      DATA A3/1.421414/
      DATA A4/-1.453152/
      DATA A5/1.061405/
      DATA P/0.327591/
      ERFNP=0.0
      IF (X.EQ.0.0) RETURN
      ES=SIGN(1.0,X)
      Y=ABS(X)
      IF (Y.LT.6.0) GO TO 1
      ERFNP=ES
      RETURN
1      Y2=Y*Y
      T=1.0/(1.0+P*Y)
      AT=((A1+(A2+(A3+(A4+A5*T)*T)*T)*T)*T)
      EAT=AT/EXP(Y2)
      ERFNP=(1.0-EAT)*ES
      RETURN
      END
      SUBROUTINE MAXLK (M,ITM,RESP,N,A,B,C,MAX,EPS,IFAIL,SDRV,THETA,TINF
10,EXPTOT,NUMITS,SEM)
      EXTERNAL FODLOG,SODLOG
      INTEGER RESP(M)
      DIMENSION A(N), B(N), C(N), ITM(M)
C*** USES MAXIMUM LIKELIHOOD LOGISTIC SCORING ALGORITHM AND RESPONSE
C*** MODEL
C*** BISECTION IS USED TO PROVIDE THE INITIAL GUESS FOR THE
C*** NEWTON-RAPHSON METHOD
      CALL BISECT (FODLOG,RESP,A,B,C,M,ITM,5,GUESS)
C
      CALL NEWTRAP (FODLOG,SODLOG,RESP,A,B,C,M,ITM,MAX,EPS,NUMITS,GUESS,
1THETA,SDRV,IFAIL)
C
      IF (IFAIL.EQ.1) 1,2
C*** NEWTON RAPHSON DID NOT CONVERGE
1      CALL NWTRP (THETA,SDRV,SEM,TINFO,EXPTOT)
      RETURN
C
2      CALL LGSTAT (M,ITM,A,B,C,THETA,TINFO,EXPTOT)
      SEM=1.0/SQRT(AHS(SDRV))
      RETURN
      END
      FUNCTION FODLOG (RESP,ITM,A,B,C,M,THETA)
      INTEGER RESP(M),RIGHT
      DIMENSION A(M), B(M), C(M), ITM(M)
      DATA XMAX,XMIN/200.0,-200.0/
      DATA D,RIGHT/1.7,1/
C*** CALCULATES FIRST DERIVATIVE OF LOG-LIKELIHOOD FUNCTION OF A
C*** RESPONSE VECTOR FOR THE LOGISTIC MODEL
      SUM=0.0
      DO 1 I=1,M
      IF (ITM(I).EQ.0) GO TO 1
      X=D*A(I)*(THETA-B(I))
      IF (X.LT.XMIN) X=XMIN
      IF (X.GT.XMAX) X=XMAX

```



```

      EXF=EXP(X)
      AE=A(I)*EXF
      SUM=SUM-AE/(EXF+1.0)
      IF (RESP(I).NE.RIGHT) GO TO 1
      CE=C(I)+EXF
      SUM=SUM+AE/CE
1     CONTINUE
      SDLOG=-1.7*SUM
      RETURN
END
FUNCTION SDLOG (RESP,ITM,A,B,C,M,THETA)
  INTEGER RESP(M),RIGHT
  DIMENSION ITM(M), A(M), B(M), C(M)
  DATA XMAX,XMIN/200.0,-200.0/
  DATA D,RIGHT/1.7,1/
C*** CALCULATES SECOND DERIVATIVE OF LOG-LIKELIHOOD FUNCTION
C*** OF A RESPONSE VECTOR FOR THE LOGISTIC MODEL
  SUM=0.0
  DO 1 I=1,M
    IF (ITM(I).EQ.0) GO TO 1
    X=D*A(I)*(THETA-B(I))
    IF (X.LT.XMIN) X=XMIN
    IF (X.GT.XMAX) X=XMAX
    EXF=EXP(X)
    AE=A(I)*EXF
    SUM=SUM-A(I)*AE/((1.0+EXF)*(1.0+EXF))
    IF (RESP(I).NE.RIGHT) GO TO 1
    CE=C(I)+EXF
    SUM=SUM+A(I)*C(I)*AE/(CE*CE)
1   CONTINUE
  SDLOG=-2.89*SUM
  RETURN
END
SUBROUTINE BISECT (F1,RESP,A,B,C,M,ITM,NITER,BMID)
  INTEGER RESP(M)
  DIMENSION A(M), B(M), C(M), ITM(M)
C*** CALCULATES APPROXIMATE ROOT OF F1 BY BISECTIONS
C*** BISECTING NITER (NUMBER OF ITERATIONS) TIMES.
C*** BMID IS BEST CURRENT GUESS AT ROOT THETA
  C
C*** INITIALIZE LEFT BOUND AND F1(BOUND) AND RIGHT BOUND F1(BOUND)
  BL=-5.0
  BR=5.0
  BMID=0.0
  FL=F1(RESP,ITM,A,B,C,M,BL)
  FR=F1(RESP,ITM,A,B,C,M,BR)
C*** TEST FOR NO ROOT IN INTERVAL--RETURN IF NO SOLUTION
  IF ((TL*TR).GT.0.0) RETURN
  C
C*** NOW CALCULATE BISECTIONS NITER TIMES
  DO 3 I=1,NITER
    TMID=F1(RESP,ITM,A,B,C,M,BMID)
    IF ((TMID*TL).GT.0.0) GO TO 1
C*** REPLACE RIGHT BOUND WITH BMID
    BR=BMID
    GO TO 2
C*** REPLACE LEFT BOUND WITH BMID
1   TL=TMID
    BL=BMID
C*** FIND NEW MIDPOINT BMID
2   BMID=(BL+BR)/2.0
3   CONTINUE
  RETURN
END
SUBROUTINE NEWTRAP (F1,F2,RESP,A,B,C,M,ITM,NITER,EPS,NUMITS,GUESS,
  1THETA,SDRV,IFAIL)

```

```

      INTEGER RESP(M)
      DIMENSION A(M), R(M), C(M), ITM(M)
C*** CALCULATES THE ROOT OF F1 GIVEN ITS FIRST DERIVATIVE F2
C*** AND AN INITIAL GUESS USING NEWTON-RAPHSON METHOD
C*** THETA IS APPR. TO THE ROOT; SORV IS F2(THETA)
      NUMITS=0
      THETA=GUESS
C*** LOOP UNTIL ERR<EPS OR NUMBER OF ITERATIONS BECOMES TOO LARGE
1   FORV=F1(RSP,ITM,A,R,C,M,THETA)
      SORV=F2(RSP,ITM,A,R,C,M,THETA)
      ERR=FORV/SORV
      THETA=THETA-ERR
      NUMITS=NUMITS+1
C*** EXIT LOOP CRITERION
      IF ((NUMITS.LT.NITER).AND.(ABS(ERR).GT.EPS)) GO TO 1
C*** END LOOP. TEST FOR CONVERGENCE AND SET IFAIL
      IFAIL=0
      IF (ABS(ERR).LT.EPS) RETURN
C
C*** NEWTON RAPHSON METHOD DOES NOT CONVERGE
      IFAIL=1
      RETURN
      END
      SUBROUTINE WHERE (THETA,SEDM,SEM,TINFO,EXPTOT)
C*** SET PARAM VALUES FOR THE CASE IN WHICH NEWTON RAPHSON FAILS
C*** TO CONVERGE
      THETA=-99.99
      SEDM=-99.99
      SEM=-99.99
      TINFO=-99.99
      EXPTOT=-99.99
      RETURN
      END
      SUBROUTINE LGSTAT (M,ITM,A,R,C,THETA,TINFO,EXPTOT)
      DIMENSION A(M), R(M), C(M), ITM(M)
      DATA XMAX,XMIN/12.0,-12.0/
      TINFO=0.0
      EXPTOT=0.0
      KOUNT=0
C
      DO 1 I=1,M
        IF (ITM(I).EQ.0) GO TO 1
        KOUNT=KOUNT+1
        ARG=-1.7*A(I)*(THETA-R(I))
        IF (ARGU.GT.XMAX) ARGU=XMAX
        IF (ARGU.LT.XMIN) ARGU=XMIN
        P=C(I)+(1.0-C(I))*(1.0/(1.0+EXP(A-60)))
        Q=1.-P
        FARG=EXP(-ARGU)
        PRIME=FARG/((1.0+FARG)*(1.0+P*Q))
        PRIME=PRIME*(1.0-C(I))*A(I)*1.7
        TINFO=TINFO+(PRIME*PRIME)/(6.*Q)
        EXPTOT=EXPTOT+P
1      CONTINUE
      EXPTOT=EXPTOT/KOUNT
      RETURN
      END
      SUBROUTINE MAXENO (RESP,ITM,M,N,A,R,C,MAX,EPS,THETA,SORV,IFAIL,TIN
      FO,EXPTOT,NUMITS,SEM)
      EXTERNAL FDNQGV,SDNQGV
      INTEGER RESP(M)
      DIMENSION ITM(N), A(N), R(N), C(N)
C*** USES MAXIMUM LIKELIHOOD NORMAL GIVE SCORING ALGORITHM AND
C*** RESPONSE VECTOR
C*** MISERION IS USED TO PROVIDE THE INITIAL GUESS FOR THE
C*** NEWTON RAPHSON METHOD

```

```

C      CALL BISECT (FONOGV,RESP,A,B,C,M,ITM,5,GUESS)
C
C      CALL NEWTRAP (FONOGV,SDNOGV,RESP,A,B,C,M,ITM,MAX,FPS,NUMITS,GUESS,
1THETA,SDRV,IFAIL)
      IF (IFAIL.EQ.1) 1,2
C*** NEWTON RAPHSON DID NOT CONVERGE
1      CALL FMTERR (THETA,SDRV,SEM,TINFO,EXPTOT)
      RETURN
C
2      CALL NOSIAT (M,ITM,A,B,C,THETA,TINFO,EXPTOT)
      SDRV=ABS(SDRV)
      SEM=1.0/SQRT(SDRV)
      RETURN
      END
      FUNCTION FONOGV (RESP,ITM,A,B,C,M,THETA)
      INTEGER RESP(M),RIGHT
      DIMENSION A(M), B(M), C(M), ITM(M)
      DATA PI,RIGHT/3.141592,1/
      DATA XMAX,XMIN/7.0,-7.0/
C*** CALCULATES FIRST DERIVATIVE OF LOG-LIKELIHOOD FUNCTION OF
C*** A RESPONSE VECTOR FOR THE NORMAL OGIVE MODEL
C
      SUM=0.0
      ROOTPI=1.0/SQRT(2.0*PI)
      DO 2 I=1,M
      IF (ITM(I).EQ.0) GO TO 2
      TEMP=A(I)*(THETA-B(I))
      IF (TEMP.GT.XMAX) TEMP=XMAX
      IF (TEMP.LT.XMIN) TEMP=XMIN
      X=-(TEMP*TEMP)/2.0
      DNMRAT=ROOTPI*A(I)*(1.0-C(I))*EXP(X)
      DENOM=C(I)+(1.0-C(I))*COFN(TEMP)
      IF (RESP(I).EQ.RIGHT) GO TO 1
      DENOM=-(1.0-DENOM)
1      SUM=SUM+(DNMRAT/DENOM)
2      CONTINUE
      FONOGV=SUM
      RETURN
      END
      FUNCTION SDNOGV (RESP,ITM,A,B,C,M,THETA)
      INTEGER RESP(M),RIGHT
      DIMENSION A(M), B(M), C(M), ITM(M)
      DATA PI,RIGHT/3.141592,1/
      DATA XMAX,XMIN/7.0,-7.0/
C*** CALCULATES SECOND DERIVATIVE OF LOG-LIKELIHOOD FUNCTION
C*** OF A RESPONSE VECTOR FOR THE NORMAL OGIVE MODEL
C
      SUM=0.0
      ROOTPI=1.0/SQRT(2.0*PI)
      DO 2 I=1,M
      IF (ITM(I).EQ.0) GO TO 2
      TEMP1=A(I)*(THETA-B(I))
      IF (TEMP1.GT.XMAX) TEMP1=XMAX
      IF (TEMP1.LT.XMIN) TEMP1=XMIN
      X=-TEMP1*TEMP1/2.0
      TEMP2=ROOTPI*(1.0-C(I))*A(I)*EXP(X)
      FIRNUM=TEMP2*TEMP2
      SECNUM=TEMP2*A(I)*TEMP1
      SDENOM=C(I)+(1.0-C(I))*COFN(TEMP1)
      FDENOM=SDENOM*SDENOM
      IF (RESP(I).EQ.RIGHT) GO TO 1
      FDENOM=-(1.0-SDENOM)*(1.0-SDENOM)
      SDENOM=-(1.0-SDENOM)
1      SUM=SUM-(FIRNUM/FDENOM)-(SECNUM/SDENOM)
2      CONTINUE

```



	SND06V=SUM	27
	RETURN	28
	END	29
	SUBROUTINE NOSTAT (M, IIM, A, R, C, THETA, TINFQ, EXPTOT)	1
	DIMENSION A(M), R(M), C(M), IIM(M)	2
	DATA XMAX, XMIN / 7.0, -7.0 /	3
	DATA PI / 3.1415927	4
	TINFQ=0.0	5
	EXPTOT=0.0	6
	KOUNT=0	7
C		8
	DO 1 I=1, M	9
	IF (IIM(I).EQ.0) GO TO 1	10
	KOUNT=KOUNT+1	11
	TEMP=A(I)*(THETA-R(I))	12
	IF (TEMP.GT.XMAX) TEMP=XMAX	13
	IF (TEMP.LT.XMIN) TEMP=XMIN	14
	P=C(I)+(1.0-C(I))*CDFN(TEMP)	15
	Q=1.-P	16
	TEMP=-TEMP*TEMP/2.0	17
	PPRIME=(1.)/SQRT(2.)*PI)*(1.-C(I))*A(I)*EXP(TEMP)	18
	TINFQ=TINFQ+(PPRIME*PPRIME)/(P*Q)	19
	EXPTOT=EXPTOT+P	20
1	CONTINUE	21
	EXPTOT=EXPTOT/KOUNT	22
	RETURN	23
	END	24

## APPENDIX E

```
PROGRAM LINPSCO (INPUT,OUTPUT,DATA,IPOOL,TAPE1=DATA,TAPE2=IPOOL,TAPE3,PUNCH)
```

C		67
C***	SEARCH FOR ITEMS ADMINISTERED IN POOL	68
	DO 9 J=1,M	69
	IKI=ADIM(NCAT(J),OPT4)	70
	K=BDIM(NCAT(J),OPT4)	71
	IF (INAD(J).NE.0) GO TO 5	72
C***	BLANK OR REJECTED ITEM ENCOUNTERED	73
	DO 3 L=1,IKI	74
3	ADM(J,L)=0.0	75
	DO 4 L=1,K	76
4	BDM(J,L)=0.0	77
	GO TO 9	78
5	CONTINUE	79
C		80
	DO 4 J=1,INUP	81
	IF (INAD(J).NE.ITEM(I)) GO TO 8	82
	DO 6 L=1,IKI	83
6	ADM(J,L)=A(I,L)	84
	DO 7 L=1,K	85
7	BDM(J,L)=B(I,L)	86
	GO TO 9	87
8	CONTINUE	88
	INAD(J)=0	89
9	CONTINUE	90
	GO TO 13	91
C		92
C***	ALL ITEMS IN POOL HAVE BEEN ADMINISTERED SO POOL DOES NOT	93
C***	NEED TO BE SEARCHED	94
10	DO 10 I=1,M	95
	IKI=ADIM(NCAT(I),OPT4)	96
	K=BDIM(NCAT(I),OPT4)	97
	DO 11 JJ=1,IKI	98
11	ADM(I,JJ)=A(I,JJ)	99
	DO 12 J=1,K	100
	BDM(I,J)=B(I,J)	101
12	CONTINUE	102
C		103
C		104
C***	PUNCH THE ITEM PARAMETERS ESTIMATES CORRESPONDING TO THE ITEMS	105
C***	IN THE TEST	106
13	IF (OPT1.NE.1) GO TO 15	107
	DO 14 I=1,M	108
	IF (INAD(I).EQ.0) GO TO 14	109
	IKI=ADIM(NCAT(I),OPT4)	110
	K=BDIM(NCAT(I),OPT4)	111
	PUNCH 35, INAD(I), IREJ(I), (ADM(I,JJ),JJ=1,IKI)	112
	PUNCH 37, (BDM(I,J),J=1,K)	113
14	CONTINUE	114
C		115
C	PRINT OUT THE ITEMS AND THEIR PARAMETERS	116
15	DO 16 III=1,M	117
	IKI=ADIM(NCAT(III),OPT4)	118
	K=BDIM(NCAT(III),OPT4)	119
	PRINT 39, INAD(III), IREJ(III), (ADM(III,JJ),JJ=1,IKI)	120
16	PRINT 40, (BDM(III,J),J=1,K)	121
	PRINT 43	122
C	*****	123
C***		124
C***	READ SUBJECT FROM TAPE1, CALCULATE THETA, LOOP BACK TO 300 UNTIL*	125
C***	THERE ARE NO MORE SUBJECTS TO SCORE	126
C***		127
C	*****	128
	N=0	129
17	READ (1,IFORM) NAME,IO, (IRESP(I),I=1,M)	130
	IO=UNIT(1)	131
	IF (IO.LE.0) GO TO 18	132



PRINT 38, IO	133
18 IF (IO.EQ.0) GO TO 29	134
C	135
C*** CHECK FOR VALID RESPONSES AND FOR SPECIAL RESPONSE VECTORS	136
CALL CHKRSR (IRES, IOMIT, INAD, NCAT, M, NAME, IO, INADS)	137
C	138
CALL CHKVEC (IRES, INADS, NCAT, M, ICHK, NBEST, NANS)	139
GO TO (19, 20, 21, 23), ICHK	140
C*** ALL ITEMS OMITTED	141
19 PRINT 45, NAME, IO	142
T=-50.00	143
PERCNT=-50.00	144
GO TO 22	145
C*** ALL RESPONSES INCORRECT	146
20 PRINT 46, NAME, IO	147
T=-10.0	148
PERCNT=0.0	149
GO TO 22	150
C*** ALL RESPONSES CORRECT	151
21 PRINT 47, NAME, IO	152
T=10.0	153
PERCNT=1.0	154
C	155
22 SFORM=0.0	156
NUMITS=0	157
TINFO=0.0	158
GO TO 28	159
23 CONTINUE	160
C	161
PERCNT=FLOAT(NBEST)/FLOAT(NANS)	162
C	163
C	164
C*** DETERMINE WHICH MODEL TO USE	165
IF (OPT4-2) 24, 25, 26	166
24 CALL LOGRAD (IRES, ADM, BDM, M, NCAT, INADS, .001, 50, SFORM, IFAIL, T, NUMI	167
ITS, TINFO, SE)	168
GO TO 27	169
25 CALL NOGRAD (IRES, ADM, BDM, M, NCAT, INADS, .001, 50, SFORM, IFAIL, T, NUMI	170
ITS, TINFO, SE)	171
GO TO 27	172
26 CALL NOMLOG (IRES, ADM, BDM, M, NCAT, INADS, .001, 50, SFORM, IFAIL, T, NUMI	173
ITS, TINFO, SE)	174
C	175
C*** OUTPUT RESULTS TO TAPE3	176
27 IF (IFAIL.EQ.0) GO TO 28	177
C*** CONVERGENCE NOT OBTAINED	178
PRINT 41, NAME, IO	179
NC=NC+1	180
C	181
28 WRITE (3, 42) NAME, IO, PERCNT, T, SFORM, NANS, NUMITS, TINFO, SE	182
N=N+1	183
GO TO 17	184
C	185
29 PRINT 44, N, NC	186
STOP	187
C	188
30 FORMAT (2I4, I1, 2I1, 1X, I1, F5.2, 27X, I2)	189
31 FORMAT (16I5)	190
32 FORMAT (80I1)	191
33 FORMAT (8A10)	192
34 FORMAT (8A10)	193
35 FORMAT (T50, *LINPSC0*, //, T50*====*, //, T20, *LINEAR POLYCHOTOMU	194
1S SCORING WITH TWO PARAMETER MODELS*, //, T40*PSYCHOMETRICS METHOD	195
2S PROGRAM*, //, T40*DEPARTMENT OF PSYCHOLOGY*, //, T40*UNIVERSITY OF MIN	196
3NESOTA*, //, T40*MPLS. MINN. 55455*, //, T20, *INUP*, T27, *=*, I4, //, T20	197
4, *MMAX*, T27, *=*I4, //, T20, *IOMIT*, T27, *=*, I4, //, T20*OPT1*, T27*==*I4, //,	198

```

51 T20*OPT2*,T27*==*,I4,/,T20*OPT4*,T27*==*,I4,/,T20,*MAXCAT*,T27,*==*,I 199
64, /T20,*U*,T27,*==*,F4.1, /T20,*VARIABLE FORMAT*,* FOR POOL =*,8A10, 200
7 /T20,*VARIABLE FORMAT FOR DATA =*,9A10, /T20,8A10, /T20,9A10, /T20,8 201
9A10) 202
36 FORMAT (15,1X,I2,2X,10F6.2) 203
37 FORMAT (10F6.2) 204
38 FORMAT (*PARITY ERROR ON TAPE*,10JX,I2) 205
39 FORMAT (/10X,*ITEM ID = *,I5,5X,*REJECTION =*,I3, /12X,*A: *,10F6. 206
12) 207
40 FORMAT (12X,*B: *,10F6.2) 208
41 FORMAT (10X,*CONVERGENCE NOT OBTAINED FOR SUBJECT =*,2A10,* ID =*, 209
1A9) 210
42 FORMAT (1X,2A10,A4,F5.2,2F7.2,I4,I4,2F7.2) 211
43 FORMAT (////) 212
44 FORMAT (10X,*CASES READ =*,I5,* CASES NOT CONVERGED =*,I5) 213
45 FORMAT (10X,*ALL ITEMS OMITTED FOR SUBJECT =*,2A10,* ID =*,A9) 214
46 FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS ALL ANSWERS IN*,*COR 215
1RECT*) 216
47 FORMAT (10X,*SUBJECT =*,2A10,* ID =*,A9,* HAS ALL ANSWERS COR*,*RE 217
1CT*) 218
END 219
INTEGER FUNCTION ADIM (NUMCAT,OPT4) 1
INTEGER OPT4 2
C*** DETERMINES THE NUMBER OF A PARAMETERS FOR A GIVEN ITEM 3
ADIM=1 4
IF (OPT4.-2.3) ADIM=NUMCAT+1 5
RETURN 6
END 7
INTEGER FUNCTION BDM (NUMCAT,OPT4) 1
INTEGER OPT4 2
C*** DETERMINES THE NUMBER OF B PARAMETERS FOR A GIVEN ITEM 3
BDM=NUMCAT 4
IF (OPT4.-2.3) BDM=NUMCAT+1 5
RETURN 6
END 7
SUBROUTINE CHKINF (INUP,M,OPT4) 1
INTEGER OPT4 2
C*** CHECKS FOR ERRORS IN THE INPUT DATA 3
C*** IF AN ERROR IS FOUND, A MESSAGE IS PRINTED AND THE PROGRAM 4
C*** HALTS 5
C 6
IERRE=0 7
IF (INUP.LE.100) GO TO 1 8
PRINT 4, INUP 9
IERRE=1 10
1 IF (M.LE.INUP) GO TO 2 11
PRINT 5, M,INUP 12
IERRE=1 13
2 IF (1.LE.OPT4.AND.OPT4.LE.3) GO TO 3 14
PRINT 6, OPT4 15
IERRE=1 16
3 IF (IERRE.EQ.1) STOP 17
RETURN 18
C 19
C 20
4 FORMAT (10X,*INPUT ERROR: NO. OF ITEMS IN ITEM POOL =*,I5, /10X,*N 21
10. MUST BE .LE. 10(*) 22
5 FORMAT (10X,*INPUT ERROR: NO. ITEMS ADMINISTERED =*,I5, /10X,*NO. 23
1MUST BE .LE. NO. OF ITEMS IN ITEM POOL =*,I5) 24
6 FORMAT (10X,*INPUT ERROR: OPTION 4 =*,I3,* DOES NOT CORRES*,*POND 25
1 TO ANY OF THE AVAILABLE RESPONSE MODELS*) 26
END 27
SUBROUTINE NOGPAD (IRESP,A,P, ITEMS,NCAT, INAD,EPS,MAXIT,SFORM,IFAI 1
1L,THE IA,NUMITS,TINFO,SE) 2
EXTERNAL FORVAD,SORVNO 3
DIMENSION IRESP(100), A(100,10), P(100,10), NCAT(10), INAD(100) 4

```

```

CALL BISECT (FORVNO, IRESP, A, B, NITEMS, NCAT, INAD, S, GUESS)
CALL NEWTAP (FORVNO, SORVNO, IRESP, A, B, NITEMS, NCAT, INAD, MAXIT, EPS, N
1UMITS, GUESS, THETA, SDRV, TFAIL)
SFORM=-SDRV
IF (IFAIL.EQ.1) GO TO 1
CALL NOINFO (A, B, NITEMS, NCAT, INAD, THETA, TINFO)
SE=1.0/SQRT(ABS(SDRV))
RETURN
C
1 TINFO=-99.99
SE=-99.99
RETURN
END
FUNCTION IVALUE (IRESP, NCAT)
C*** CHECKS RESPONSE FOR SPECIAL CASES
C*** IVALUE RETURNS . . . 1 IF IRESP IS BEST RESPONSE
C*** 2 IF IRESP IS WORST RESPONSE
C*** 3 OTHERWISE
C
IVALUE=3
IF (IRESP.EQ.1) IVALUE=1
IF (IRESP.EQ.(NCAT+1)) IVALUE=2
RETURN
END
SUBROUTINE RCAT1 (A, B, THETA, TMINB0, TMINB1, P, ETOZ0, ETOZ1)
C*** COMPUTES VALUES NECESSARY FOR THE CALCULATION OF THE DERIV-
C*** ATIVES OF THE NORMAL OGIVE GRADED MODEL FOR THE SPECIAL CASE
C*** WHEN IRESP IS THE BEST RESPONSE
C
TMINB0=THETA-B
TMINB1=0.0
Y=A*TMINB0
P=CDFN(Y)
ETOZ0=EXP(-Y*Y/2.0)
ETOZ1=0.0
RETURN
END
SUBROUTINE RCATN (A, B, THETA, TMINB0, TMINB1, P, ETOZ0, ETOZ1)
C*** COMPUTES VALUES NECESSARY FOR THE CALCULATION OF THE DERIV-
C*** ATIVES OF THE NORMAL OGIVE GRADED MODEL FOR THE SPECIAL
C*** CASE WHEN IRESP IS THE WORST RESPONSE
C
TMINB0=0.0
TMINB1=THETA-B
Y1=A*TMINB1
P=1.0-CDFN(Y1)
ETOZ0=0.0
ETOZ1=EXP(-Y1*Y1/2.0)
RETURN
END
SUBROUTINE RCATOT (A, B0, B1, THETA, TMINB0, TMINB1, P, ETOZ0, ETOZ1)
C*** COMPUTES VALUES NECESSARY FOR THE CALCULATION OF DERIVATIVES OF
C*** THE NORMAL OGIVE GRADED MODEL FOR ALL OTHER CASES
C
TMINB0=THETA-B0
TMINB1=THETA-B1
Y=A*TMINB0
Y1=A*TMINB1
P=CDFN(Y)-CDFN(Y1)
ETOZ0=EXP(-Y*Y/2.0)
ETOZ1=EXP(-Y1*Y1/2.0)
RETURN
END
FUNCTION FORVNO (IRESP, INAD, NITEMS, NCAT, A, B, THETA)
C*** DIMENSION IRESP(100), INAD(100), NCAT(100), A(100,10), B(100,10)
C*** CALCULATES THE FIRST DERIVATIVE FOR THE NORMAL OGIVE GRADED MODEL

```



```

C
SUM=0.0
DO 5 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 5
K=I-RESP(I)
J=IVALUE(K,NCAT(I))
GO TO (1,2,3), J
C
C*** IRESP IS BEST RESPONSE
1 CALL FCA11 (A(I,1),B(I,K),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
C*** IRESP IS WORST RESPONSE
2 CALL FCA11 (A(I,1),B(I,K-1),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
C*** ALL OTHER RESPONSES
3 CALL FCA11 (A(I,1),B(I,K),B(I,K-1),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
IF (P.EQ.0.0) P=0.0001
SUM=SUM+A(I,1)*(ETOZ0-ETOZ1)/P
CONTINUE
C
SOVNO=SUM/ROOT(2.0*3.142)
RETURN
END
FUNCTION SOVNO (IRESP,INAD,NITEMS,NCAT,A,B,THETA)
DIMENSION IRESP(100), INAD(100), NCAT(100), A(100,10), B(100,10)
C*** CALCULATE SECOND DERIVATIVE FOR THE NORMAL OGIVE GRADED MODEL
C
SUM=0.0
ROOTPI=1.0/ROOT(2.0*3.142)
DO 5 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 5
K=I-RESP(I)
J=IVALUE(K,NCAT(I))
GO TO (1,2,3), J
C
C*** IRESP IS BEST RESPONSE
1 CALL FCA11 (A(I,1),B(I,K),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
C*** IRESP IS WORST RESPONSE
2 CALL FCA11 (A(I,1),B(I,K-1),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
C*** ALL OTHER RESPONSES
3 CALL FCA11 (A(I,1),B(I,K),B(I,K-1),THETA,TMINB0,TMINB1,P,ETOZ0,ETOZ1)
GO TO 4
C
IF (P.EQ.0.0) P=0.0001
SUM1=A(I,1)*ROOTPI*(ETOZ0-ETOZ1)/P
SUM2=-SUM1*SUM1
SUM3=-(A(I,1)**3)*ROOTPI*((TMINB0*ETOZ0)-(TMINB1*ETOZ1))/P
SUM=SUM+SUM1+SUM2
CONTINUE
C
SOVNO=SUM
RETURN
END
SUBROUTINE FCA11 (A,B,NITEMS,NCAT,IAD,THETA,TINFO)
DIMENSION A(100,10), B(100,10), NCAT(100), IAD(100)
C*** COMPUTES INFORMATION FOR GRADED NORMAL OGIVE MODEL AT GIVEN

```

```

C*** VALUE OF THETA
C
TINFO=0.0
ROOTPI=1.0/SQRT(2.0*3.142)
C*** LOOP OVER ITEMS
DO 5 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 5
C
C*** INITIALIZATION--VALUES CALCULATED FOR FIRST CATEGORY OF ITEM I
Y=A(I,1)*(THETA-B(I,1))
P0=CHFN(Y)
P1=0.0
ETOZ0=EXP(-Y*Y/2.0)
ETOZ1=0.0
KATGRY=0
C
C*** LOOP OVER ALL THE CATEGORIES OF ITEM I
1 KATGRY=KATGRY+1
P=P0-P1
IF (P.EQ.0.0) P=0.0001
FDRVP=A(I,1)*ROOTPI*(ETOZ0-ETOZ1)
TINFO=TINFO+(FDRVP*FDRVP)/P
C
ETOZ1=ETOZ0
P1=P0
C
IF (KATGRY-(NCAT(I)+1)) 2,3,4
C*** CURRENT CATEGORY UNDER CONSIDERATION IS NOT ONE OF THE EXTREMES
2 Y=A(I,1)*(THETA-B(I,KATGRY))
P0=CHFN(Y)
ETOZ0=EXP(-Y*Y/2.0)
GO TO 1
C*** LAST CATEGORY OF AN ITEM IS BEING CONSIDERED
3 P0=1.0
ETOZ0=0.0
GO TO 1
C*** ALL CATEGORIES FOR ITEM I HAVE BEEN EXAMINED
4 CONTINUE
C
5 CONTINUE
RETURN
END
SUBROUTINE LOGRAD (IRESP,A,B,NITEMS,NCAT,INAD,EPS,NITER,SFORM,IFAI
1L,THETA,NUMITS,TINFO,SE)
EXTERNAL FDRVLL,SORVLL
DIMENSION IRESP(100), A(100,10), B(100,10), NCAT(100), INAD(100)
CALL BISECT (FDRVLL,IRESP,A,B,NITEMS,NCAT,INAD,5,GUESS)
CALL NEWTRAP (FDRVLL,SORVLL,IRESP,A,B,NITEMS,NCAT,INAD,NITER,EPS,N
1UMITS,GUESS,THETA,SORV,IFAIL)
SFORM=-SORV
IF (IFAIL.EQ.1) GO TO 1
CALL LLINFO (A,B,NITEMS,NCAT,INAD,THETA,TINFO)
SE=1.0/SQRT(ARS(SORV))
RETURN
C
1 TINFO=-99.99
SE=-99.99
RETURN
END
SUBROUTINE CALCP (IRESP,INAD,NCAT,A,B,NITEMS,THETA,P)
COMMON D
DIMENSION IRESP(100), INAD(100), NCAT(100), B(100,10), P(100,2), A
1(100,10)
C*** CALCULATES UPPER AND LOWER P FOR EACH ITEM WITH GIVEN ANSWER VECT
DO 4 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 4

```

```

J=IRESP(I)
IF (J.EQ.1) GO TO 1
P(I,1)=1.0/(1.0+EXP(-D*A(I,1)*(THETA-B(I,J-1))))
GO TO 2
1 P(I,1)=0.0
2 IF (J.EQ.(NCAT(I)+1)) GO TO 3
P(I,2)=1.0/(1.0+EXP(-D*A(I,1)*(THETA-B(I,J))))
GO TO 4
3 P(I,2)=1.0
4 CONTINUE
RETURN
END
FUNCTION FORVLL (IRESP,INAD,NITEMS,NCAT,A,R,THETA)
COMMON D
DIMENSION P(100,2), A(100,10), R(100,10), INAD(100), IRESP(100), N
1 NCAT(100)
C*** CALCULATES FIRST DERIVATIVE OF LOG-LIKE FUNCTION
SUM=0.0
CALL CALCF (IRESP,INAD,NCAT,A,R,NITEMS,THETA,P)
DO 1 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 1
SUM=SUM+A(I,1)*(1.0-P(I,1)-P(I,2))
1 CONTINUE
FORVLL=SUM*D
RETURN
END
FUNCTION SDPVLL (IRESP,INAD,NITEMS,NCAT,A,R,THETA)
COMMON D
DIMENSION P(100,2), A(100,10), R(100,10), INAD(100), IRESP(100), N
1 NCAT(100)
C*** CALCULATES SECOND DERIVATIVE OF LOGLIKE FUNCTION
SUM=0.0
CALL CALCF (IRESP,INAD,NCAT,A,R,NITEMS,THETA,P)
DO 1 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 1
Q1=1.0-P(I,1)
Q2=1.0-P(I,2)
P1=P(I,2)-P(I,1)
DIFF1=2.0*Q1-1.0
DIFF2=2.0*Q2-1.0
R1=P(I,1)*Q1
R2=P(I,2)*Q2
SUM=SUM+(A(I,1)**2/P1)*(-DIFF1*R1+DIFF2*R2-((R1-R2)**2)/P1)
1 CONTINUE
SDPVLL=D*D*SUM
RETURN
END
SUBROUTINE PCAT (A,R,THETA,ITEM,NUMCAT,FLOWER,PUPPER)
COMMON D
DIMENSION B(100,10), FLOWER(10), PUPPER(10)
C*** CALCULATES P'S FOR ALL RESPONSE CATEGORIES OF A GIVEN ITEM
C
FLOWER(1)=0.0
DO 1 I=1,NUMCAT
PUPPER(I)=1.0/(1.0+EXP(-D*A*(THETA-B(ITEM,I))))
FLOWER(I+1)=PUPPER(I)
1 CONTINUE
PUPPER(NUMCAT+1)=1.0
RETURN
END
SUBROUTINE LLINFO (A,R,NITEMS,NCAT,INAD,THETA,TINFO)
COMMON D
DIMENSION A(100,10), R(100,10), NCAT(100), INAD(100)
DIMENSION PUPPER(10), FLOWER(10)
C*** COMPUTES INFORMATION FOR THE GRADED LOGISTIC MODEL AT A GIVEN
C*** VALUE OF THETA

```



```

C
  TINFO=0.0
  DO 1 I=1,NITEMS
    IF (INAD(I).EQ.0) GO TO 2
    CALL PCAT (A(I,1),B,THETA,I,NCAT(I),PLOWER,PUPPER)
  1
C*** LOOP OVER ALL THE RESPONSE CATEGORIES OF ITEM I
  NCATIG=NCAT(I)+1
  DO 1 J=1,NCATEG
    QUPPER=1-PUPPER(J)
    QLOWER=1-PLOWER(J)
    P=PUPPER(J)-PLOWER(J)
    IF (P.EQ.0.0) P=0.0001
    FDRVP=0*A(I,1)*(PUPPER(J)*QUPPER-PLOWER(J)*QLOWER)
    TINFO=TINFO+(FDRV*FDRVP)/P
  1 CONTINUE
  2 CONTINUE
  RETURN
END
SUBROUTINE NOMLOG (IRESP,A,B,NITEMS,NCAT,INAD,EPS,MAXIT,SFORM,IFAI
1L,THETA,NUMITS,TINFO,SE)
  EXTERNAL FDRVNL
  DIMENSION IRESP(100), A(100,10), B(100,10), NCAT(100), INAD(100)
  CALL MISECT (FDRVNL,IRESP,A,B,NITEMS,NCAT,INAD,5,GUESHI,GUESLO)
  CALL SECANT (FDRVNL,IRESP,A,B,NITEMS,NCAT,INAD,MAXIT,EPS,GUESHI,GU
1ESLO,NUMITS,THETA,SDRV,IFAIL)
  SFORM=SDRV
  IF (IFAIL.EQ.1) GO TO 1
  CALL NLINEF (A,B,NITEMS,NCAT,INAD,THETA,TINFO)
  SE=1.0/SQRT(ABS(SDRV))
  RETURN
C
1 TINFO=-99.99
  SE=-99.99
  RETURN
END
FUNCTION FDRVNL (IRESP,INAD,NITEMS,NCAT,A,B,THETA)
  DIMENSION IRESP(100), INAD(100), NCAT(100), A(100,10), B(100,10)
C*** CALCULATES FIRST DERIVATIVE OF NOMINAL LOGISTIC FUNCTION
C*** NOTE: FOR THIS MODEL, THE A'S ARE THE SLOPE PARAMETERS
C*** AND THE B'S ARE THE INTERCEPT PARAMETERS
C
  SUM=0.0
  DO 2 I=1,NITEMS
    IF (INAD(I).EQ.0) GO TO 2
    XNUM=0.0
    DENOM=0.0
    NUMCAT=NCAT(I)+1
    DO 1 J=1,NUMCAT
      CONST=A(I,IRESP(I))-A(I,J)
      ARG=A(I,J)+THETA*B(I,J)
      EZ=EXP(ARG)
      XNUM=XNUM+CONST*EZ
      DENOM=DENOM+EZ
    1 CONTINUE
    SUM=SUM+XNUM/DENOM
  2 CONTINUE
  FDRVNL=SUM
  RETURN
END
SUBROUTINE NLINEF (A,B,NITEMS,NCAT,INAD,THETA,TINFO)
  DIMENSION A(100,10), B(100,10), NCAT(100), INAD(100)
  DIMENSION E(10)
C*** COMPUTES INFORMATION FOR THE NOMINAL LOGISTIC FUNCTION AT A
C*** GIVEN VALUE OF THETA
C

```

```

      INFO=0.0
C*** LOOP OVER THE ITEMS
      DO 4 I=1,NITEMS
      IF (INAD(I).EQ.0) GO TO 4
      NCATG=NCAT(I)+1
C
C*** COMPUTE THE DENOMINATOR USED TO CALCULATE P AND THE FIRST
C*** DERIVATIVE OF P--IT IS THE SAME VALUE FOR ALL POSSIBLE RE-
C*** SPONSES TO ITEM I
      DENOM=0.0
      DO 1 K=1,NCATEG
      E(K)=EXP(A(I,K)*THETA+B(I,K))
      DENOM=DENOM+E(K)
1      CONTINUE
C
C*** SUM OVER ALL POSSIBLE RESPONSES
      DO 3 J=1,NCATEG
      P=E(J)/DENOM
C
C*** CALCULATE THE FIRST DERIVATIVE OF P
      DENVNUM=0.0
      DO 2 K=1,NCATEG
      DENVNUM=DENVNUM+E(K)*(A(I,J)-A(I,K))
2      CONTINUE
      FDRVP=(E(J)*DENVNUM)/(DENOM*DENOM)
C
      INFO=INFO+(FDRVP*FDRVP)/P
3      CONTINUE
4      CONTINUE
      RETURN
      END
      SUBROUTINE CHKRSR (IRESP,IOMIT,INAD,NCAT,NITEMS,NAME,ID,INADS)
      DIMENSION IRESP(100), INAD(100), NCAT(100), NAME(2), INADS(100)
C*** DELETES OMITTED ITEMS FOR A SUBJECT FROM LIST OF ITEMS
C*** ADMINISTERED (RESULTS RETURNED IN INADS) AND CHECKS TO SEE
C*** IF ALL RESPONSES ARE VALID. IF NOT AN ERROR MESSAGE IS WRITTEN
C*** AND THE PROGRAM HALTS
C
      DO 2 I=1,NITEMS
      NUMCAT=NCAT(I)+1
      INADS(I)=INAD(I)
      IF (IRESP(I).NE.IOMIT) GO TO 1
      INADS(I)=0
      GO TO 2
C
C*** CHECK FOR INVALID RESPONSE
1      IF ((1.LE.IRESP(I)).AND.(IRESP(I).LE.NUMCAT)) GO TO 2
      PRINT 3, NAME,ID,IRESP(I),NUMCAT
      STOP
C
2      CONTINUE
      RETURN
C
3      FORMAT (10X,'ERROR IN DATA: SUBJECT = ',2A10,' ID = ',49,' HAS IL-
      LEGAL RESPONSE =',I3,'/10X,' RESPONSES MUST BE',* BETWEEN 1 AND NUMC-
      TER OF CATEGORIES =',I3,' INCLUSIVE')
      END
      SUBROUTINE CHKVEC (IRESP,INAD,NCAT,NITEMS,ICLK,NBEST,NANSI)
      DIMENSION IRESP(100), INAD(100), NCAT(100)
C
C*** CHKVEC TESTS THE RAW RESPONSE VECTOR TO DETERMINE IF
C IF IT IS ONE OF THE FOLLOWING THREE CASES:
C*** ALL BEST RESPONSES...ICLK SET TO 3
C*** ALL WORST RESPONSES...ICLK SET TO 2
C*** NO RESPONSES.....ICLK SET TO 1
C*** OTHERWISE.....ICLK SET TO 4

```

```

NANS=0
NWORST=0
NREST=0
DO 1 I=1,NITEMS
IF (INAD(I).EQ.0) GO TO 1
NANS=NANS+1
IF (IRESP(I).EQ.1) NREST=NREST+1
IF (IRESP(I).EQ.NCAT(I)+1) NWORST=NWORST+1
1 CONTINUE
C
ICHK=4
IF (NANS.EQ.0) ICHK=1
IF (NWORST.EQ.NANS) ICHK=2
IF (NREST.EQ.NANS) ICHK=3
RETURN
END
SUBROUTINE BISECT (F1,IRESP,A,B,NITEMS,NCAT,INAD,NIL,BMID,BLAST)
DIMENSION IRESP(100), A(100,10), B(100,10), NCAT(100), INAD(100),
1P(100,2)
C*** CALCULATES APPROXIMATE ROOT OF F1 BY BISECTION,BISECTING THE INTER
C*** NIT (NUMBER OF ITERATIONS) TIMES. BMID IS BEST CURRENT GUESS AT T
C*** BLAST IS THE PREVIOUS VALUE OF BMID. IT IS USED AS THE SECOND
C*** INITIAL GUESS FOR THE SECANT METHOD.
C
C INITIALIZE LEFT BOUND AND F1(BOUND), AND RIGHT BOUND,F1(BOUND)
BL=-5.0
BR=5.0
BMID=0.0
TL=F1(IRESP,INAD,NITEMS,NCAT,A,B,BL)
TR=F1(IRESP,INAD,NITEMS,NCAT,A,B,BR)
C TEST FOR NO ROOT IN INTERVAL, AND RETURN IF NOT POSSIBLE
IF ((IL*TR).GT.0.0) RETURN
C NOW CALCULATE BISECTIONS NIT TIMES
DO 3 I=1,NIT
TMID=F1(IRESP,INAD,NITEMS,NCAT,A,B,BMID)
IF ((TMID*TL).GT.0.0) GO TO 1
BR=BMID
GO TO 2
1 TL=TMID
2 BL=BMID
3 BMID=(BL+BR)/2.0
CONTINUE
BLAST=BR
IF (IL.EQ.TMID) BLAST=BL
RETURN
END
SUBROUTINE NEWTRAP (F1,F2,IRESP,A,B,NITEMS,NCAT,INAD,NITER,EPS,NUM
1ITS,GUESS,THETA,SORV,IFAIL)
DIMENSION IRESP(100), A(100,10), B(100,10), NCAT(100), INAD(100)
C*** CALCULATES ROOT OF F1 GIVEN ITS FIRST DERIVATIVE F2 AND AN INITIAL
C*** GUESS. UTILIZES NEWTON-RAPHSONS METHOD
C
C INITIALIZE
NUMITS=0
THETA=GUESS
C*** LOOP UNTIL ERR IS LESS THAN EPS OR NUMBER OF ITERATIONS BECOMES TO
1 SORV=F2(IRESP,INAD,NITEMS,NCAT,A,B,THETA)
FORV=F1(IRESP,INAD,NITEMS,NCAT,A,B,THETA)
ERR=FORV/SORV
THETA=THETA-ERR
NUMITS=NUMITS+1
C EXIT LOOP CRITERION
IF ((NUMITS.LT.NITER).AND.(ABS(ERR).GT.EPS)) GO TO 1
C*** END LOOP, TEST FOR FAILURE AND SET IFAIL
IFAIL=0
IF (ABS(ERR).LT.EPS) RETURN
20

```



```

IFAIL=1
THETA=-99.99
SDRV=99.99
RETURN
END
SUBROUTINE SECANT (F1,IRESP,A,B,NITEMS,NCAT,INAD,MAXIT,EPS,GUESHI,
1 GUESLO,NUMITS,THETA,SLOPE,IFAIL)
2 DIMENSION IRESP(100), A(100,10), B(100,10), NCAT(100), INAD(100)
3
C*** USES THE SECANT METHOD TO CALCULATE THE ROOT, THETA, OF THE
C*** FUNCTION F1
C*** GUESHI AND GUESLO ARE THE TWO INITIAL GUESSES AT THE ROOT
C*** REQUIRED BY THE SECANT METHOD
C
4 NUMITS=0
5 THETA=GUESHI
6 FLAST=GUESLO
7 FLAST=F1(IRESP,INAD,NITEMS,NCAT,A,B,FLAST)
8
C
9
10
11
12
13
C*** LOOP UNTIL CONVERGENCE OR NONCONVERGENCE IS ESTABLISHED
14
1 FCUR=F1(IRESP,INAD,NITEMS,NCAT,A,B,THETA)
15
16 IF (FCUR.EQ.FLAST) GO TO 2
17 SLOPE=(THETA-FLAST)/(FCUR-FLAST)
18 CHANGE=FCUR*SLOPE
19 FLAST=THETA
20 FLAST=FCUR
21 THETA=THETA-CHANGE
22 NUMITS=NUMITS+1
23 IF (ABS(CHANGE).GT.EPS.AND.NUMITS.LT.MAXIT) GO TO 1
24
C
25 IFAIL=0
26 SLOPE=1.0/SLOPE
27 IF (ABS(CHANGE).LT.EPS) RETURN
28
C*** SECANT METHOD DOES NOT CONVERGE IN MAXIT ITERATIONS
29 IFAIL=1
30 THETA=-99.99
31 SLOPE=-99.99
32 RETURN
33
C
34
C*** ERROR: SECANT METHOD CANNOT BE USED ON F1
35
2 PRINT 3
36 IFAIL=1
37 THETA=-99.99
38 SLOPE=-99.99
39 RETURN
40
C
41
3
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252
2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
2289
2290
2291
2292
2293
2294
2295
2296
2297
2298
2299
2300
2301
2302
2303
2304
2305
2306
2307
2308
2309
2310
2311
2312
2313
2314
2315
2316
2317
2318
2319
2320
2321
2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334
2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375
2376
2377
2378
2379
2380
2381
2382
2383
2384
2385
2386
2387
2388
2389
2390
2391
2392
2393
2394
2395
2396
2397
2398
2399
2400
2401
2402
2403
2404
2405
2406
2407
2408
2409
2410
2411
2412
2413
2414
2415
2416
2417
2418
2419
2420
2421
2422
2423
2424
2425
2426
2427
2428
2429
2430
2431
2432
2433
2434
2435
2436
2437
2438
2439
2440
2441
2442
2443
2444
2445
2446
2447
2448
2449
2450
2451
2452
2453
2454
2455
2456
2457
2458
2459
2460
2461
2462
2463
2464
2465
2466
2467
2468
2469
2470
2471
2472
2473
2474
2475
2476
2477
2478
2479
2480
2481
2482
2483
2484
2485
2486
2487
2488
2489
2490
2491
2492
2493
2494
2495
2496
2497
2498
2499
2500
2501
2502
2503
2504
2505
2506
2507
2508
2509
2510
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520
2521
2522
2523
2524
2525
2526
2527
2528
2529
2530
2531
2532
2533
2534
2535
2536
2537
2538
2539
2540
2541
2542
2543
2544
2545
2546
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595

```

# DISTRIBUTION LIST

Navy		1 Scientific Advisor to the Chief of Naval Personnel (Pers-Or) Naval Bureau of Personnel Room 4410, Arlington Annex Washington, DC 20370
1 Dr. Ed Aiken Navy Personnel R&D Center San Diego, CA 92152	1 Commanding Officer Naval Health Research Center Attn: Library San Diego, CA 92152	1 DR. RICHARD A. POLLAK ACADEMIC COMPUTING CENTER U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402
1 Dr. Jack R. Forsting Provost & Academic Dean U.S. Naval Postgraduate School Monterey, CA 93940	1 Naval Medical R&D Command Code 44 National Naval Medical Center Bethesda, MD 20014	1 Mr. Arnold Rubenstein Naval Personnel Support Technology Naval Material Command (08T244) Room 1044, Crystal Plaza #5 2221 Jefferson Davis Highway Arlington, VA 20360
1 Dr. Robert Breaux Code M-71 NAVTREAEQUIPCEN Orlando, FL 32813	1 Library Navy Personnel R&D Center San Diego, CA 92152	1 A. A. SJOHOLM TECH. SUPPORT, CODE 201 NAVY PERSONNEL R&D CENTER SAN DIEGO, CA 92152
1 MR. MAURICE CALLAHAN Pers 23a Bureau of Naval Personnel Washington, DC 20370	6 Commanding Officer Naval Research Laboratory Code 2627 Washington, DC 20390	1 Mr. Robert Smith Office of Chief of Naval Operations OP-987E Washington, DC 20350
1 DR. PAT FEDERICO NAVY PERSONNEL R&D CENTER SAN DIEGO, CA 92152	1 OFFICE OF CIVILIAN PERSONNEL (CODE 26) DEPT. OF THE NAVY WASHINGTON, DC 20390	1 Dr. Alfred F. Smode Training Analysis & Evaluation Group (TAEG) Dept. of the Navy Orlando, FL 32813
1 Dr. Paul Foley Navy Personnel R&D Center San Diego, CA 92152	1 JOHN OLSEN CHIEF OF NAVAL EDUCATION & TRAINING SUPPORT PENSACOLA, FL 32509	1 Dr. Richard Sorensen Navy Personnel R&D Center San Diego, CA 92152
1 Dr. John Ford Navy Personnel R&D Center San Diego, CA 92152	1 Psychologist ONR Branch Office 495 Summer Street Boston, MA 02210	1 CDR Charles J. Theisen, JR. MSC, USN Head Human Factors Engineering Div. Naval Air Development Center Warminster, PA 18974
1 CAPT. D.M. GRAGG, MC, USN HEAD, SECTION ON MEDICAL EDUCATION UNIFORMED SERVICES UNIV. OF THE HEALTH SCIENCES 6917 ARLINGTON ROAD BETHESDA, MD 20014	1 Psychologist ONR Branch Office 536 S. Clark Street Chicago, IL 60605	1 W. Gary Thomson Naval Ocean Systems Center Code 7132 San Diego, CA 92152
1 Dr. Norman J. Kerr Chief of Naval Technical Training Naval Air Station Memphis (75) Millington, TN 38054	1 Code 436 Office of Naval Research Arlington, VA 22217	1 Dr. Ronald Weitzman Department of Administrative Sciences U. S. Naval Postgraduate School Monterey, CA 93940
1 Dr. Leonard Kroeker Navy Personnel R&D Center San Diego, CA 92152	1 Office of Naval Research Code 437 800 N. Quincy Street Arlington, VA 22217	1 DR. MARTIN F. WISKOFF NAVY PERSONNEL R&D CENTER SAN DIEGO, CA 92152
1 CHAIRMAN, LEADERSHIP & LAW DEPT. DIV. OF PROFESSIONAL DEVELOPMENT U.S. NAVAL ACADEMY ANNAPOLIS, MD 21402	5 Personnel & Training Research Program (Code 458) Office of Naval Research Arlington, VA 22217	
1 Dr. William L. Maloy Principal Civilian Advisor for Education and Training Naval Training Command, Code 00A Pensacola, FL 32508	1 Psychologist OFFICE OF NAVAL RESEARCH BRANCH 223 OLD MARYLEBONE ROAD LONDON, NW, 15TH ENGLAND	Army
1 CAPT Richard L. Martin USS Francis Marion (LPA-249) FPO New York, NY 09501	1 Psychologist ONR Branch Office 1030 East Green Street Pasadena, CA 91101	1 Technical Director U. S. Army Research Institute for the Behavioral and Social Sciences 5001 Eisenhower Avenue Alexandria, VA 22333
1 Dr. James McBride Code 301 Navy Personnel R&D Center San Diego, CA 92152	1 Scientific Director Office of Naval Research Scientific Liaison Group/Tokyo American Embassy APO San Francisco, CA 96503	1 HQ USAREUE & 7th Army ODCSOPS USAREUE Director of GED APO New York 09403
2 Dr. James McGrath Navy Personnel R&D Center Code 306 San Diego, CA 92152	1 Head, Research, Development, and Studies (OP102X) Office of the Chief of Naval Operations Washington, DC 20370	1 DR. RALPH CANTER U.S. ARMY RESEARCH INSTITUTE 5001 EISENHOWER AVENUE ALEXANDRIA, VA 22333
1 DR. WILLIAM MONTAGUE LRDC UNIVERSITY OF PITTSBURGH 3939 O'HARA STREET PITTSBURGH, PA 15213		

1 DR. RALPH DUSEK U.S. ARMY RESEARCH INSTITUTE 5001 EISENHOWER AVENUE ALEXANDRIA, VA 22333	1 Dr. Roger Pennell AFHRL/TT Lowry AFB, CO 80230	1 Military Assistant for Training and Personnel Technology Office of the Under Secretary of Defense for Research & Engineering Room 3D129, The Pentagon Washington, DC 20301
1 Dr. Myron Fischl U.S. Army Research Institute for the Social and Behavioral Sciences 5001 Eisenhower Avenue Alexandria, VA 22333	1 Personnel Analysis Division HQ USAF/DPXXA Washington, DC 20330	1 MAJOR Wayne Sellman, USAF Office of the Assistant Secretary of Defense (MRA&L) 3B930 The Pentagon Washington, DC 20301
1 Dr. Ed Johnson Army Research Institute 5001 Eisenhower Blvd. Alexandria, VA 22333	1 Research Branch AFMPC/DPMYP Randolph AFB, TX 78148	
1 Dr. Michael Kaplan U.S. ARMY RESEARCH INSTITUTE 5001 EISENHOWER AVENUE ALEXANDRIA, VA 22333	1 Dr. Malcolm Ree AFHRL/PED Brooks AFB, TX 78235	Civil Govt
1 Dr. Milton S. Katz Individual Training & Skill Evaluation Technical Area U.S. Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333	1 Dr. Marty Rockway (AFHRL/TT) Lowry AFB Colorado 80230	1 Dr. Susan Chipman Basic Skills Program National Institute of Education 1200 19th Street NW Washington, DC 20208
1 Dr. Harold F. O'Neill, Jr. ATTN: PERL-QK 5001 EISENHOWER AVENUE ALEXANDRIA, VA 22333	1 Jack A. Thorpe, Capt, USAF Program Manager Life Sciences Directorate AFOSR Holling AFB, DC 20332	1 Dr. William Gorham, Director Personnel R&D Center U.S. Civil Service Commission 1900 E Street NW Washington, DC 20415
1 Dr. Robert Ross U.S. Army Research Institute for the Social and Behavioral Sciences 5001 Eisenhower Avenue Alexandria, VA 22333	1 Brian K. Waters, LCOL, USAF Air University Maxwell AFB Montgomery, AL 36112	1 Dr. Joseph I. Lipson Division of Science Education Room W-638 National Science Foundation Washington, DC 20550
1 Director, Training Development U.S. Army Administration Center ATTN: Dr. Sherrill Ft. Benjamin Harrison, IN 46218	Marines	
1 Dr. Frederick Steinheiser U. S. Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333	1 Director, Office of Manpower Utilization HQ, Marine Corps (MPU) BCE, Bldg. 2009 Quantico, VA 22134	1 Dr. John Mays National Institute of Education 1200 19th Street NW Washington, DC 20208
1 Dr. Joseph Ward U.S. Army Research Institute 5001 Eisenhower Avenue Alexandria, VA 22333	1 MCDEC Quantico Marine Corps Base Quantico, VA 22134	1 Dr. Arthur Melmed National Institute of Education 1200 19th Street NW Washington, DC 20208
Air Force	1 DR. A.L. SLAFKOSKY SCIENTIFIC ADVISOR (CODE RD-1) HQ, U.S. MARINE CORPS WASHINGTON, DC 20380	1 Dr. Andrew R. Molnar Science Education Dev. and Research National Science Foundation Washington, DC 20550
1 Air Force Human Resources Lab AFHRL/PED Brooks AFB, TX 78235	CoastGuard	
1 Air University Library AUL/LSE 76/443 Maxwell AFB, AL 36112	1 MR. JOSEPH J. COWAN, CHIEF PSYCHOLOGICAL RESEARCH (G-P-1/62) U.S. COAST GUARD HQ WASHINGTON, DC 20590	1 Dr. Lalitha P. Sanathanan Environmental Impact Studies Division Argonne National Laboratory 9700 S. Cass Avenue Argonne, IL 60439
1 Dr. Philip De Leo AFHRL/II Lowry AFB, CO 80230	1 Dr. Thomas Warm U. S. Coast Guard Institute P. O. Substation 18 Oklahoma City, OK 73169	1 Dr. Jeffrey Schiller National Institute of Education 1200 19th St. NW Washington, DC 20208
1 DR. G. A. ECKSTRAND AFHRL/AS WRIGHT-PATTERSON AFB, OH 45433	Other DoD	1 Dr. Thomas G. Sticht Basic Skills Program National Institute of Education 1200 19th Street NW Washington, DC 20208
1 CDR. MERCER CNET LIAISON OFFICER AFHRL/FLYING TRAINING DIV. WILLIAMS AFB, AZ 85224	12 Defense Documentation Center Cameron Station, Bldg. 5 Alexandria, VA 22314 Attn: TC	1 Dr. Vern W. Urry Personnel R&D Center U.S. Civil Service Commission 1900 E Street NW Washington, DC 20415
1 Dr. Ross L. Morgan (AFHRL/ASR) Wright -Patterson AFB Ohio 45433	1 Dr. Dexter Fletcher ADVANCED RESEARCH PROJECTS AGENCY 1400 WILSON BLVD. ARLINGTON, VA 22209	1 Dr. Joseph L. Young, Director Memory & Cognitive Processes National Science Foundation Washington, DC 20550



# Non Govt

- 1 Dr. Earl A. Allvusi  
HQ, AFHRL (AFSC)  
Brooks AFB, TX 78235
- 1 Dr. Erling E. Anderson  
University of Copenhagen  
Studiestraedt  
Copenhagen  
DENMARK
- 1 1 psychological research unit  
Dept. of Defense (Army Office)  
Campbell Park Offices  
Canberra ACT 2600, Australia
- 1 Dr. Alan Haddley  
Medical Research Council  
Applied Psychology Unit  
15 Chaucer Road  
Cambridge CB2 2EF  
ENGLAND
- 1 Dr. Isaac Bejar  
Educational Testing Service  
Princeton, NJ 08450
- 1 Dr. Warner Birice  
Streitkraefteamt  
Rosenberg 5300  
Eonn, West Germany D-5300
- 1 Dr. R. Darrel Bock  
Department of Education  
University of Chicago  
Chicago, IL 60637
- 1 Dr. Nicholas A. Bond  
Dept. of Psychology  
Sacramento State College  
600 Jay Street  
Sacramento, CA 95819
- 1 Dr. David G. Powers  
Institute for Social Research  
University of Michigan  
Ann Arbor, MI 48106
- 1 Dr. Robert Brennan  
American College Testing Progr  
P. O. Box 168  
Iowa City, IA 52240
- 1 DR. C. VICTOR BUNDERSON  
WICAT INC.  
UNIVERSITY PLAZA, SUITE 10  
1160 SO. STATE ST.  
OREM, UT 84057
- 1 Dr. John B. Carroll  
Psychometric Lab  
Univ. of No. Carolina  
Davie Hall 013A  
Chapel Hill, NC 27514
- 1 Charles Myers Library  
Livingstone House  
Livingstone Road  
Stratford  
London E15 2LJ  
ENGLAND
- 1 Dr. Kenneth E. Clark  
College of Arts & Sciences  
University of Rochester  
River Campus Station  
Rochester, NY 14627
- 1 Dr. Norman Cliff  
Dept. of Psychology  
Univ. of So. California  
University Park  
Los Angeles, CA 90007
- 1 Dr. William Coffman  
Iowa Testing Programs  
University of Iowa  
Iowa City, IA 52242
- 1 Dr. Allan M. Collins  
Bolt Beranek & Newman, Inc.  
50 Moulton Street  
Cambridge, Ma 02138
- 1 Dr. Meredith Crawford  
Department of Engineering Administration  
George Washington University  
Suite 805  
2101 L Street N. W.  
Washington, DC 20037
- 1 Dr. Hans Cronbag  
Education Research Center  
University of Leyden  
Boerhaavelaan 2  
Leyden  
The NETHERLANDS
- 1 MAJOR I. N. EVONIC  
CANADIAN FORCES PERS. APPLIED RESEARCH  
1107 AVENUE ROAD  
TORONTO, ONTARIO, CANADA
- 1 Dr. Leonard Feldt  
Lindquist Center for Measurment  
University of Iowa  
Iowa City, IA 52242
- 1 Dr. Richard L. Ferguson  
The American College Testing Program  
P.O. Box 168  
Iowa City, IA 52240
- 1 Dr. Victor Fields  
Dept. of Psychology  
Montgomery College  
Rockville, MD 20850
- 1 Dr. Gerhardt Fischer  
Liebigasse 5  
Vienna 1010  
Austria
- 1 Dr. Donald Fitzgerald  
University of New England  
Armidale, New South Wales 2351  
AUSTRALIA
- 1 Dr. Edwin A. Fleishman  
Advanced Research Resources Organ.  
Suite 900  
4330 East West Highway  
Washington, DC 20014
- 1 Dr. John R. Frederiksen  
Bolt Beranek & Newman  
50 Moulton Street  
Cambridge, MA 02138
- 1 DR. ROBERT GLASER  
LRDC  
UNIVERSITY OF PITTSBURGH  
3939 O'HARA STREET  
PITTSBURGH, PA 15213
- 1 Dr. Ross Greene  
CTB/McGraw Hill  
Del Monte Research Park  
Monterey, CA 93940
- 1 Dr. Alan Cross  
Center for Advanced Study in Education  
City University of New York  
New York, NY 10026
- 1 Dr. Ron Hambleton  
School of Education  
University of Massachusetts  
Amherst, MA 01002
- 1 Dr. Chester Harris  
School of Education  
University of California  
Santa Barbara, CA 93106
- 1 Dr. Lloyd Humphreys  
Department of Psychology  
University of Illinois  
Champaign, IL 61820
- 1 Library  
HumRRC/Western Division  
27857 Herwick Drive  
Carmel, CA 93821
- 1 Dr. Steven Hunka  
Department of Education  
University of Alberta  
Edmonton, Alberta  
CANADA
- 1 Dr. Earl Hunt  
Dept. of Psychology  
University of Washington  
Seattle, WA 98105
- 1 Dr. Huynh Huynh  
Department of Education  
University of South Carolina  
Columbia, SC 29208
- 1 Dr. Carl J. Jensema  
Gallaudet College  
Kendall Green  
Washington, DC 20002
- 1 Dr. Arnold F. Kanarick  
Honeywell, Inc.  
2600 Ridgeway Pkwy  
Minneapolis, MN 55413
- 1 Dr. John A. Keats  
University of Newcastle  
Newcastle, New South Wales  
AUSTRALIA
- 1 Mr. Marlin Kroger  
1117 Via Coleta  
Palos Verdes Estates, CA 90274
- 1 LCOL. C.R.J. LAFLEUR  
PERSONNEL APPLIED RESEARCH  
NATIONAL DEFENSE HQS  
101 COLONEL BY DRIVE  
OTTAWA, CANADA K1A 0K2
- 1 Dr. Michael Levine  
Department of Psychology  
University of Illinois  
Champaign, IL 61820
- 1 Dr. Robert Linn  
College of Education  
University of Illinois  
Urbana, IL 61801
- 1 Dr. Frederick M. Lord  
Educational Testing Service  
Princeton, NJ 08540

- 1 Dr. Robert R. Mackie  
Human Factors Research, Inc.  
6780 Cortona Drive  
Santa Barbara Research Pk.  
Goleta, CA 93017
- 1 Dr. Gary Marco  
Educational Testing Service  
Princeton, NJ 08450
- 1 Dr. Scott Maxwell  
Department of Psychology  
University of Houston  
Houston, TX 77025
- 1 Dr. Sam Mayo  
Loyola University of Chicago  
Chicago, IL 60601
- 1 Dr. Allen Munro  
Univ. of So. California  
Behavioral Technology Labs  
3717 South Hope Street  
Los Angeles, CA 90007
- 1 Dr. Melvin R. Novick  
Iowa Testing Programs  
University of Iowa  
Iowa City, IA 52242
- 1 Dr. Jesse Orlansky  
Institute for Defense Analysis  
400 Army Navy Drive  
Arlington, VA 22202
- 1 Dr. James A. Paulson  
Portland State University  
P.O. Box 751  
Portland, OR 97207
- 1 MR. LUIGI PETRULLO  
2431 N. EDGEWOOD STREET  
ARLINGTON, VA 22207
- 1 DR. STEVEN M. PINE  
4050 Douglas Avenue  
Golden Valley, MN 55416
- 1 DR. DIANE M. RAMSEY-KLEE  
R-K RESEARCH & SYSTEM DESIGN  
3947 RIDGEMONT DRIVE  
MALIBU, CA 90265
- 1 MIN. RET. M. RAUCH  
P II 4  
BUNDESMINISTERIUM DER VERTEIDIGUNG  
POSTFACH 161  
53 BONN 1, GERMANY
- 1 Dr. Peter F. Read  
Social Science Research Council  
605 Third Avenue  
New York, NY 10016
- 1 Dr. Mark D. Reckase  
Educational Psychology Dept.  
University of Missouri-Columbia  
12 Hill Hall  
Columbia, MO 65201
- 1 Dr. Fred Reif  
SESAME  
c/o Physics Department  
University of California  
Berkeley, CA 94720
- 1 Dr. Andrew M. Rose  
American Institutes for Research  
1055 Thomas Jefferson St. NW  
Washington, DC 20007
- 1 Dr. Leonard L. Rosenbaum, Chairman  
Department of Psychology  
Montgomery College  
Rockville, MD 20850
- 1 Dr. Ernst Z. Rothkopf  
Bell Laboratories  
600 Mountain Avenue  
Murray Hill, NJ 07974
- 1 Dr. Donald Rubin  
Educational Testing Service  
Princeton, NJ 08450
- 1 Dr. Larry Rudner  
Gallaudet College  
Kendall Green  
Washington, DC 20002
- 1 Dr. J. Ryan  
Department of Education  
University of South Carolina  
Columbia, SC 29208
- 1 PROF. FUMIKO SAMEJIMA  
DEPT. OF PSYCHOLOGY  
UNIVERSITY OF TENNESSEE  
KNOXVILLE, TN 37916
- 1 DR. ROBERT J. SEIDEL  
INSTRUCTIONAL TECHNOLOGY GROUP  
HUMPRO  
300 N. WASHINGTON ST.  
ALEXANDRIA, VA 22314
- 1 Dr. Kazuo Shigemasa  
University of Tohoku  
Department of Educational Psychology  
Kawauchi, Sendai 982  
JAPAN
- 1 Dr. Edwin Shirkey  
Department of Psychology  
Florida Technological University  
Orlando, FL 32816
- 1 Dr. Richard Snow  
School of Education  
Stanford University  
Stanford, CA 94305
- 1 Dr. Robert Sternberg  
Dept. of Psychology  
Yale University  
Box 11A, Yale Station  
New Haven, CT 06520
- 1 DR. ALBERT STEVENS  
BOLT BERANEK & NEWMAN, INC.  
50 MOULTON STREET  
CAMBRIDGE, MA 02138
- 1 DR. PATRICK SUPPES  
INSTITUTE FOR MATHEMATICAL STUDIES IN  
THE SOCIAL SCIENCES  
STANFORD UNIVERSITY  
STANFORD, CA 94305
- 1 Dr. Hariharan Swaminathan  
Laboratory of Psychometric and  
Evaluation Research  
School of Education  
University of Massachusetts  
Amherst, MA 01003
- 1 Dr. Brad Sympson  
Elliott Hall  
University of Minnesota  
75 E. River Road  
Minneapolis, MN 55455
- 1 Dr. Kikumi Tatsuoka  
Computer Based Education Research  
Laboratory  
252 Engineering Research Laboratory  
University of Illinois  
Urbana, IL 61801
- 1 Dr. David Thissen  
Department of Psychology  
University of Kansas  
Lawrence, KS 66044
- 1 Dr. J. Uhlaner  
Perceptronics, Inc.  
6271 Variel Avenue  
Woodland Hills, CA 91364
- 1 Dr. Howard Wainer  
Bureau of Social Science Research  
1990 M Street, N. W.  
Washington, DC 20036
- 1 DR. THOMAS WALLSTEN  
PSYCHOMETRIC LABORATORY  
DAVIE HALL 013A  
UNIVERSITY OF NORTH CAROL  
CHAPEL HILL, NC 27514
- 1 Dr. John Wannous  
Department of Management  
Michigan University  
East Lansing, MI 48824
- 1 DR. SUSAN E. WHITELEY  
PSYCHOLOGY DEPARTMENT  
UNIVERSITY OF KANSAS  
LAWRENCE, KANSAS 66044
- 1 Dr. Wolfgang Wildgrube  
Streitkrafteamt  
Rosenberg 5300  
Bonn, West Germany D-5300
- 1 Dr. Robert Woud  
School Examination Department  
University of London  
56-72 Gower Street  
London WC1E 6EE  
ENGLAND
- 1 Dr. Karl Zinn  
Center for research on Learning  
and Teaching  
University of Michigan  
Ann Arbor, MI 48104